

+15

1. A sphere of radius R carries polarization $\vec{P}(\vec{r}) = kr^2\hat{r}$.

(a) Calculate the bound charges ρ_b and σ_b .

(b) Find the electric field \vec{E} inside and outside the sphere. Check that the appropriate boundary conditions are satisfied.

+20

2. A solid sphere with radius R has uniform polarization $\vec{P} = P\hat{z}$. Also at radius R , there is a uniform surface charge density σ_f .

(a) Determine the bound charge distribution.

(b) Find the electric field \vec{E} everywhere.

(c) Find the field \vec{D} everywhere.

(d) Check that the \vec{E} and \vec{D} fields satisfy the proper boundary conditions at the surface.

(Determine ΔE_r , ΔE_θ , ΔD_r and ΔD_θ at the surface.)

Helpful information for Problem 2

A spherical shell with surface charge $\sigma(\theta) = \sigma_0 \cos \theta$ and radius R has scalar potential

$$V(r, \theta) = \frac{\sigma_0}{3\epsilon_0} r \cos \theta, \quad r \leq R$$

$$= \frac{\sigma_0}{3\epsilon_0} \frac{R^3}{r^2} \cos \theta, \quad r \geq R$$

+20

3. Consider an infinite solenoid with a central infinite wire oriented along its axis. The central wire carries steady current $I_z\hat{z}$. The solenoid (radius R) carries steady current $I_\phi\hat{\phi}$ and has n turns per unit length.

(a) Find the magnetic field \vec{B} everywhere. What are the boundary conditions on \vec{B} at the surface of the solenoid?

(b) Find the vector potential \vec{A} everywhere. What are the boundary conditions on \vec{A} at the surface of the solenoid?

+15 4. A long solenoid is filled with a magnetic material with uniform magnetization \vec{M} aligned with the solenoid axis. The solenoid has n turns per unit length, and it has radius R . A steady current I flows through the solenoid coils.

- Find the magnetic field \vec{B} .
- Find \vec{H} everywhere.
- How are \vec{B} and \vec{H} related if the magnetized material inside the solenoid is a linear medium?

+15 5. Two infinite planes parallel to the xy -plane carry steady surface currents $\vec{K}(t) = \pm \hat{x} K_0 \cos \omega t$, respectively. A finite solenoid with center axis aligned along the \hat{y} axis is placed between the planes, without any current. The solenoid has resistance R and radius a . Find the current induced in the solenoid as a function of time.

length l
 n turns/length

+15 6. Consider two point charges q at $z = a$ and $-q$ at $z = -a$. Determine the force of the point charge $-q$ on the point charge $+q$ by integrating Maxwell's stress tensor dotted into the infinitesimal area element $d\vec{a}$ over the xy plane. Recall that Maxwell's stress tensor is defined by

$$T_{ij} \equiv \epsilon_0 \left(E_i E_j - \frac{1}{2} E^2 \right) + \frac{1}{\mu_0} \left(B_i B_j - \frac{1}{2} B^2 \right)$$

δ_{ij} δ_{ij}

VECTOR DERIVATIVES

Cartesian. $d\mathbf{l} = dx\hat{\mathbf{x}} + dy\hat{\mathbf{y}} + dz\hat{\mathbf{z}}$; $d\tau = dx\,dy\,dz$

$$\text{Gradient : } \nabla r = \frac{\partial r}{\partial x}\hat{\mathbf{x}} + \frac{\partial r}{\partial y}\hat{\mathbf{y}} + \frac{\partial r}{\partial z}\hat{\mathbf{z}}$$

$$\text{Divergence : } \nabla \cdot \mathbf{v} = \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z}$$

$$\text{Curl : } \nabla \times \mathbf{v} = \left(\frac{\partial v_z}{\partial y} - \frac{\partial v_y}{\partial z} \right) \hat{\mathbf{x}} + \left(\frac{\partial v_x}{\partial z} - \frac{\partial v_z}{\partial x} \right) \hat{\mathbf{y}} + \left(\frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \right) \hat{\mathbf{z}}$$

$$\text{Laplacian : } \nabla^2 r = \frac{\partial^2 r}{\partial x^2} + \frac{\partial^2 r}{\partial y^2} + \frac{\partial^2 r}{\partial z^2}$$

Spherical. $d\mathbf{l} = dr\hat{\mathbf{r}} + r d\theta\hat{\theta} + r \sin\theta d\phi\hat{\phi}$; $d\tau = r^2 \sin\theta dr d\theta d\phi$

$$\text{Gradient : } \nabla r = \frac{\partial r}{\partial r}\hat{\mathbf{r}} + \frac{1}{r} \frac{\partial r}{\partial \theta}\hat{\theta} + \frac{1}{r \sin\theta} \frac{\partial r}{\partial \phi}\hat{\phi}$$

$$\text{Divergence : } \nabla \cdot \mathbf{v} = \frac{1}{r^2} \frac{\partial}{\partial r} (r^2 v_r) + \frac{1}{r \sin\theta} \frac{\partial}{\partial \theta} (\sin\theta v_\theta) + \frac{1}{r \sin\theta} \frac{\partial v_\phi}{\partial \phi}$$

$$\text{Curl : } \nabla \times \mathbf{v} = \frac{1}{r \sin\theta} \left[\frac{\partial}{\partial \theta} (\sin\theta v_\phi) - \frac{\partial v_\theta}{\partial \phi} \right] \hat{\mathbf{r}}$$

$$+ \frac{1}{r} \left[\frac{1}{\sin\theta} \frac{\partial v_r}{\partial \phi} - \frac{\partial}{\partial r} (r v_\phi) \right] \hat{\theta} + \frac{1}{r} \left[\frac{\partial}{\partial r} (r v_\theta) - \frac{\partial v_r}{\partial \theta} \right] \hat{\phi}$$

$$\text{Laplacian : } \nabla^2 r = \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial r}{\partial r} \right) + \frac{1}{r^2 \sin\theta} \frac{\partial}{\partial \theta} \left(\sin\theta \frac{\partial r}{\partial \theta} \right) + \frac{1}{r^2 \sin^2\theta} \frac{\partial^2 r}{\partial \phi^2}$$

Cylindrical. $d\mathbf{l} = ds\hat{\mathbf{s}} + s d\phi\hat{\phi} + dz\hat{\mathbf{z}}$; $d\tau = s ds d\phi dz$

$$\text{Gradient : } \nabla r = \frac{\partial r}{\partial s}\hat{\mathbf{s}} + \frac{1}{s} \frac{\partial r}{\partial \phi}\hat{\phi} + \frac{\partial r}{\partial z}\hat{\mathbf{z}}$$

$$\text{Divergence : } \nabla \cdot \mathbf{v} = \frac{1}{s} \frac{\partial}{\partial s} (sv_s) + \frac{1}{s} \frac{\partial v_\phi}{\partial \phi} + \frac{\partial v_z}{\partial z}$$

$$\text{Curl : } \nabla \times \mathbf{v} = \left[\frac{1}{s} \frac{\partial v_z}{\partial \phi} - \frac{\partial v_\phi}{\partial z} \right] \hat{\mathbf{s}} + \left[\frac{\partial v_s}{\partial z} - \frac{\partial v_z}{\partial s} \right] \hat{\phi} + \frac{1}{s} \left[\frac{\partial}{\partial s} (sv_\phi) - \frac{\partial v_s}{\partial \phi} \right] \hat{\mathbf{z}}$$

$$\text{Laplacian : } \nabla^2 r = \frac{1}{s} \frac{\partial}{\partial s} \left(s \frac{\partial r}{\partial s} \right) + \frac{1}{s^2} \frac{\partial^2 r}{\partial \phi^2} + \frac{\partial^2 r}{\partial z^2}$$

VECTOR IDENTITIES

Triple Products

$$(1) \quad \mathbf{A} \cdot (\mathbf{B} \times \mathbf{C}) = \mathbf{B} \cdot (\mathbf{C} \times \mathbf{A}) = \mathbf{C} \cdot (\mathbf{A} \times \mathbf{B})$$

$$(2) \quad \mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = \mathbf{B}(\mathbf{A} \cdot \mathbf{C}) - \mathbf{C}(\mathbf{A} \cdot \mathbf{B})$$

Product Rules

$$(3) \quad \nabla(fg) = f(\nabla g) + g(\nabla f)$$

$$(4) \quad \nabla(\mathbf{A} \cdot \mathbf{B}) = \mathbf{A} \times (\nabla \times \mathbf{B}) + \mathbf{B} \times (\nabla \times \mathbf{A}) + (\mathbf{A} \cdot \nabla)\mathbf{B} + (\mathbf{B} \cdot \nabla)\mathbf{A}$$

$$(5) \quad \nabla \cdot (f\mathbf{A}) = f(\nabla \cdot \mathbf{A}) + \mathbf{A} \cdot (\nabla f)$$

$$(6) \quad \nabla \cdot (\mathbf{A} \times \mathbf{B}) = \mathbf{B} \cdot (\nabla \times \mathbf{A}) - \mathbf{A} \cdot (\nabla \times \mathbf{B})$$

$$(7) \quad \nabla \times (f\mathbf{A}) = f(\nabla \times \mathbf{A}) - \mathbf{A} \times (\nabla f)$$

$$(8) \quad \nabla \times (\mathbf{A} \times \mathbf{B}) = (\mathbf{B} \cdot \nabla)\mathbf{A} - (\mathbf{A} \cdot \nabla)\mathbf{B} + \mathbf{A}(\nabla \cdot \mathbf{B}) - \mathbf{B}(\nabla \cdot \mathbf{A})$$

Second Derivatives

$$(9) \quad \nabla \cdot (\nabla \times \mathbf{A}) = 0$$

$$(10) \quad \nabla \times (\nabla f) = 0$$

$$(11) \quad \nabla \times (\nabla \times \mathbf{A}) = \nabla(\nabla \cdot \mathbf{A}) - \nabla^2 \mathbf{A}$$

FUNDAMENTAL THEOREMS

$$\text{Gradient Theorem : } \int_{\mathbf{a}}^{\mathbf{b}} (\nabla f) \cdot d\mathbf{l} = f(\mathbf{b}) - f(\mathbf{a})$$

$$\text{Divergence Theorem : } \int (\nabla \cdot \mathbf{A}) d\tau = \oint \mathbf{A} \cdot d\mathbf{s}$$

$$\text{Curl Theorem : } \int (\nabla \times \mathbf{A}) \cdot d\mathbf{s} = \oint \mathbf{A} \cdot d\mathbf{l}$$

BASIC EQUATIONS OF ELECTRODYNAMICS

FUNDAMENTAL CONSTANTS

Maxwell's Equations

In general :

$$\left\{ \begin{array}{l} \nabla \cdot \mathbf{E} = \frac{1}{\epsilon_0} \rho \\ \nabla \times \mathbf{E} = - \frac{\partial \mathbf{B}}{\partial t} \\ \nabla \times \mathbf{B} = \mu_0 \mathbf{J} + \mu_0 \epsilon_0 \frac{\partial \mathbf{E}}{\partial t} \\ \nabla \cdot \mathbf{B} = 0 \end{array} \right.$$

In matter :

$$\left\{ \begin{array}{l} \nabla \cdot \mathbf{D} = \rho_f \\ \nabla \times \mathbf{E} = - \frac{\partial \mathbf{B}}{\partial t} \\ \nabla \cdot \mathbf{B} = 0 \end{array} \right.$$

SPHERICAL AND CYLINDRICAL COORDINATES

Spherical

$$\left\{ \begin{array}{l} x = r \sin \theta \cos \phi \\ y = r \sin \theta \sin \phi \\ z = r \cos \theta \end{array} \right.$$

Cylindrical

$$\left\{ \begin{array}{l} r = \sqrt{x^2 + y^2 + z^2} \\ \theta = \tan^{-1}(\sqrt{x^2 + y^2}/z) \\ \phi = \tan^{-1}(y/x) \end{array} \right.$$

Cylindrical

$$\left\{ \begin{array}{l} x = s \cos \phi \\ y = s \sin \phi \\ z = z \end{array} \right.$$

Cylindrical

$$\left\{ \begin{array}{l} s = \sqrt{x^2 + y^2} \\ \phi = \tan^{-1}(y/x) \\ z = z \end{array} \right.$$

Energy, Momentum, and Power

Energy :
$$U = \frac{1}{2} \int \left(\epsilon_0 E^2 + \frac{1}{\mu_0} B^2 \right) d\tau$$

Momentum :
$$\mathbf{P} = \epsilon_0 \int (\mathbf{E} \times \mathbf{B}) d\tau$$

Poynting vector :
$$\mathbf{S} = \frac{1}{\mu_0} (\mathbf{E} \times \mathbf{B})$$

Larmor formula :
$$P = \frac{\mu_0}{6\pi c} q^2 a^2$$

$\epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{Nm}^2$

(permittivity of free space)

$\mu_0 = 4\pi \times 10^{-7} \text{ N/A}^2$

(permeability of free space)

$c = 3.00 \times 10^8 \text{ m/s}$

(speed of light)

$e = 1.60 \times 10^{-19} \text{ C}$

(charge of the electron)

$m = 9.11 \times 10^{-31} \text{ kg}$

(mass of the electron)