

Physics 214 UCSD/225a UCSB

Lecture 8

- Finish neutrino Physics
 - Neutrinos going through Matter
 - Large Mixing MSW effect
 - Using Matter effect to understand the mass hierarchy.
 - Summary of everything we know
 - Are neutrinos their own anti-particles ?
 - Nuclear double beta decay
- Aside on number of light neutrinos from LEP.

References for Neutrino Physics

- B.Kayser hep-ph/0506165
 - Most of what I've done comes from there.
- 3 NuSAG reports to HEPAP
 - 1st for Majorana neutrinos
 - 2nd for $\sin \theta_{13}$
 - 3rd for where the field is going.
 - Have used that the least in these lectures.
- All of this is linked into the course web page.

Neutrinos going through Matter

- Elastic scattering in the forward direction modifies the wave propagation by introducing something akin to an index of refraction. $e^{ipL} \rightarrow e^{inpL}$

$$n = 1 + \frac{2\pi N_e f(0)}{p^2}$$

- In EWK theory, one can show that:
 - Effect has opposite sign for neutrinos and anti-neutrinos.
 - $N = 1 - \sqrt{2} G_F N_e / p$
- We will not go through the derivation of what this does to the oscillation amplitude. I refer you to Kayser's paper for that. Instead, I'll simply quote the result and then discuss the impact on nature.

Impact of Matter on Oscillation

Oscillation probability in vacuum:

$$\text{Prob}(\nu_e \rightarrow \nu_\mu) = \sin^2 2\theta \left[\sin^2 \frac{(m_1^2 - m_2^2)L}{4E} \right]$$

Oscillation probability in matter:

$$\text{Prob}(\nu_e \rightarrow \nu_\mu) = \sin^2 2\theta_M \left[\sin^2 \frac{\Delta m_M^2 L}{4E} \right]$$

$$\sin^2 2\theta_M = \frac{\sin^2 2\theta}{\sin^2 2\theta + (\cos 2\theta - x)^2}$$

$$\Delta m_M^2 = \Delta m^2 \sqrt{\sin^2 2\theta + (\cos 2\theta - x)^2}$$

$$x = \frac{2\sqrt{2}G_F N_e E}{\Delta m^2}$$

Plug in some numbers:

- Example $L=1000\text{km}$ and atmospheric neutrino oscillation:

$$|x| \sim E/12\text{GeV}$$

\Rightarrow Effect sizeable for neutrino energy $>10\text{GeV}$.

- Example solar neutrino oscillation

- $\sqrt{2} G_F N_e \sim 0.75 \cdot 10^{-5} \text{ eV}^2/\text{MeV}$


$$|x| \sim E/5\text{MeV}$$

\Rightarrow Effect sizeable for neutrinos from ${}^8\text{B}$ but not for neutrinos from ${}^7\text{Be}$ or pp.


$$|\chi| \sim E/5\text{MeV}$$

Importance of matter effect depends on the part of the spectrum an experiment is sensitive to.

For ${}^8\text{B}$ neutrinos matter effect completely dominates!



Two Possible Mass Hierarchies



Determining the Mass hierarchy from matter effect

- Measure appearance of electrons and positrons with two different beams:


$$\nu_{\mu} \rightarrow \nu_e \text{ and } \bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$$

- For beams of $E < 2\text{GeV}$, we can approximate:

$$\sin^2 2\theta_M = \sin^2 2\theta_{13} \left[1 \pm S \frac{E}{6\text{GeV}} \right]$$

$$\frac{P(\nu_{\mu} \rightarrow \nu_e)}{P(\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e)} = \begin{array}{l} >1 \text{ for } S = +, \text{ i.e. "normal"} \\ <1 \text{ for } S = -, \text{ i.e. "inverted"} \end{array}$$

- Here S is the sign of Δm_{32}



$\nu_e [|U_{ei}|^2]$


$\nu_\mu [|U_{\mu i}|^2]$

$\nu_\tau [|U_{\tau i}|^2]$

General Arguments


- L/E defines the mass splitting an experiment is sensitive to, i.e. either “atmospheric” or “solar” split.
- Pick a neutrino flavor for your source, and check if that flavor oscillates at that mass splitting.
 - If it doesn’t then one of the two states has no component for that flavor.
 - If it does, then both states have some component of that flavor.
 - If it does maximally mix then the two flavors it mixes are equally present in both states.
- Solar neutrino is special for ${}^8\text{B}$ because it measures the ν_e content of ν_2 directly.
- Unitarity then demands that ν_1 has large ν_e component to compensate for ν_3 .

Are neutrinos their own anti-particles ?



For inverted, the m_3 part is negligible.

For normal, all three can contribute equally, and this phases may render $m_{\beta\beta}$ to be arbitrarily close to zero.



Summary - the case for $\sin 2\theta_{13}$

- We have learned that resolution of all the important open questions revolve around $\sin 2\theta_{13}$, in some sense:
 - Our ability to resolve the hierarchy of neutrino masses from matter effects is aided by a large $\sin 2\theta_{13}$
 - Size of CP violation depends directly on $\sin \theta_{13}$
 - Whether or not we will conclusively show that neutrinos are NOT their own anti-particles depends on the hierarchy and thus $\sin 2\theta_{13}$.

Let's take a closer look at measurement strategies.

Towards determining $\sin\theta_{13}$

- Two strategies:
 - Electron anti-neutrino disappearance at reactors

$$P[\bar{\nu}_e \rightarrow \text{Not } \bar{\nu}_e] \cong \sin^2 2\theta_{13} \sin^2 \Delta_{31} + \cos^4 \theta_{13} \sin^2 2\theta_{12} \sin^2 \Delta_{21}$$

- (Anti-)Electron-neutrino appearance in long baseline (anti-)muon-neutrino beams.

$$\begin{aligned} P[\bar{\nu}_\mu \rightarrow \bar{\nu}_e] &\cong \sin^2 2\theta_{13} \sin^2 \theta_{23} \sin^2 \Delta_{31} \\ &+ \sin 2\theta_{13} \cos \theta_{13} \sin 2\theta_{23} \sin 2\theta_{12} \sin \Delta_{31} \sin \Delta_{21} \cos(\Delta_{32} \pm \delta) \\ &+ \sin^2 2\theta_{12} \cos^2 \theta_{23} \cos^2 \theta_{13} \sin^2 \Delta_{21} \end{aligned}$$

Reactor anti-neutrino disappearance:

$$P[\bar{\nu}_e \rightarrow \text{Not } \bar{\nu}_e] \cong \sin^2 2\theta_{13} \sin^2 \Delta_{31} + \cos^4 \theta_{13} \sin^2 2\theta_{12} \sin^2 \Delta_{21}$$

For $\sin^2 2\theta_{13} > 0.01$, the first term dominates near $|\Delta_{31}| = \pi/2$.

***Given a MINOS precision of 10% on Δm^2_{31}
This approach could yield an unambiguous result
as long as $\sin^2(2\theta_{13})$ is not too small.***

Nova and off-axis muon neutrinos.

$$\begin{aligned} P[\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e] &\cong \sin^2 2\theta_{13} \sin^2 \theta_{23} \sin^2 \Delta_{31} \\ &+ \sin 2\theta_{13} \cos \theta_{13} \sin 2\theta_{23} \sin 2\theta_{12} \sin \Delta_{31} \sin \Delta_{21} \cos(\Delta_{32} \pm \delta) \\ &+ \sin^2 2\theta_{12} \cos^2 \theta_{23} \cos^2 \theta_{13} \sin^2 \Delta_{21} \end{aligned}$$

Disappearance will get you some resolution of this mess:

$$P[\nu_{\mu} \rightarrow \text{Not } \nu_{\mu}] \cong \sin^2 2\theta_{23} \sin^2 \Delta_{atm}$$

In addition, you'd clearly want the reactor program come back with an unambiguous measurement of $\sin^2(2\theta_{13})$.

And then there is neutrino
astrophysics ...

Which I will not be talking about !
(Amusing factoid, there's a supernova
warning system worldwide.)

Number of light neutrino families.

- LEP studied $e^+ e^-$ sitting near the Z resonance.

- They measured:

- Γ_{total} the total Z width

- M_Z the Z mass

- σ_{peak} the cross section at the peak.

$$\sigma_{\text{peak}} = \frac{12\pi}{M_Z^2} \frac{\Gamma_{ee} \Gamma_f}{\Gamma_Z^2} (1 - \delta_{\text{rad}})$$


- They take from theory:

- Γ_{ee} , Γ_{hadrons} , $\Gamma_{\nu\nu}$

- They then use: $\Gamma_{\text{total}} = 3\Gamma_{ee} + \Gamma_{\text{hadrons}} + N_\nu \Gamma_{\nu\nu}$ to obtain the number of neutrino families.

- (Z. Phys. C62 (1994) 539-550)

Truth in advertizing:
the details on how
this was done for the
result shown here are
slightly different from
what was done in the
first ALEPH paper.
Previous page describes
the early not the final paper.



Basic ideas are the same.

$$N_\nu = 2.983 \pm 0.034$$