

Formulas:

$$\vec{F}_2 = \frac{q_1 q_2}{r_{21}^2} \hat{r}_{21} \quad \text{Coulomb's law ; } \vec{E} = \vec{F}/q_0 \quad \text{electric field ; } \vec{E}(x,y,z) = \int \frac{\rho(x',y',z')(\hat{r} - \hat{r}')}{|\vec{r} - \vec{r}'|^2} dx' dy' dz'$$

$$\oint \vec{E} \cdot d\vec{a} = 4\pi q_{enc} = 4\pi \int \rho dv \quad \text{Gauss' law} \quad \text{1 charge at the origin : } \vec{E}(\vec{r}) = \frac{q}{r^2} \hat{r}$$

Linear, surface, volume charge density : $dq = \lambda ds$, $dq = \sigma dA$, $dq = \rho dV$

Electric field of : charge : $E = \frac{q}{r^2}$; line of charge : $E = \frac{2\lambda}{r}$; sheet of charge : $E = 2\pi\sigma$

Potential of single charge q : $\phi(\vec{r}) = \frac{q}{r}$; charge distribution : $\phi(\vec{r}) = \int \frac{\rho(\vec{r}')}{|\vec{r} - \vec{r}'|} dx' dy' dz'$


$$\phi(x,y,z) - \phi(x_0,y_0,z_0) = - \int_{(x_0,y_0,z_0)}^{(x,y,z)} \vec{E} \cdot d\vec{s} \quad ; \quad \vec{E} = -\nabla\phi \quad ; \quad \nabla^2\phi = -4\pi\rho \quad ; \quad \text{div}\vec{E} = 4\pi\rho$$

$$\text{div}\vec{E} = \frac{\partial E_x}{\partial x} + \frac{\partial E_y}{\partial y} + \frac{\partial E_z}{\partial z} \quad ; \quad \nabla^2 = \frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} + \frac{\partial^2}{\partial z^2}$$

energy of 3 charges : $U = \frac{q_1 q_2}{r_{12}} + \frac{q_2 q_3}{r_{23}} + \frac{q_1 q_3}{r_{13}}$; energy of q in potential ϕ : $U = q \phi(x,y,z)$

Problem 1 (10 pts)

Consider the 3 charges along the x-axis at distance a from each other of same magnitude and sign as shown in the figure:



assume the center charge is at $x=0$.

- (a) Make a plot of the potential $\phi(x)$ versus x extending from large negative x to large positive x .
 - (b) Locate 2 points in the graph where if you put a test charge q_0 it will be in equilibrium. Will the equilibrium be stable or unstable? Answer separately for $q_0 > 0$ and $q_0 < 0$. (assume the test charge can only move along the x -axis.)
 - (c) Locate a point on the x axis to which you can bring a test charge q_0 from infinity without doing any net work.
- Justify all your answers.

Problem 2 (10 pts)

An infinitely long cylinder of radius R has a non-uniform charge distribution ρ in its interior. The potential for $r < R$ (r is the distance to the cylinder axis) is given by

$$\phi(x, y, z) = x^4 + 2x^2y^2 + y^4$$


(a) Find the charge density $\rho(x, y, z)$ inside the cylinder. Show that it can be expressed as $\rho(r)$.

(b) Find the electric field at $r=R$ using the charge density found in (a) and Gauss' law. Show all the steps.

Hint: $\int dx dy \cdot f(\sqrt{x^2 + y^2}) = 2\pi \int dr \cdot r \cdot f(r)$

(c) Find the electric field $\vec{E}(x = R, y = 0, z = 0)$ directly from the potential ϕ . Explain why your result agrees or disagrees with the result in (b).

Problem 3 (10 pts)



Consider a ring of radius a and total charge q , i.e. linear charge density $\lambda = q / (2\pi a)$.

(a) Find the potential at point P a distance z along the axis from the center.

(b) Calculate the electric field (in the z direction) at point P from the potential. Make a plot of the electric field E_z versus z that includes both positive and negative z .

(c) Calculate the electric field directly from its definition, without using the potential.

Explain all steps. Explain why your result does or does not agree with the result of (b).