

## 7.2 Wave Nature of Matter

De Broglie Wavelength  
 Diffraction of electrons  
 Uncertainty Principle  
 Wave Function  
 Tunneling

## Wave properties of matter

Material particles behave as waves with a wavelength given by the De Broglie wavelength (Planck's constant/momentum)

$$\lambda = \frac{h}{p}$$

The particles are diffracted by passing through an aperture in a similar manner as light waves.

The wave properties of particles mean that when you confine it in a very small space its momentum (and kinetic energy) must increase. (uncertainty principle) This is responsible for the size of the atom.

Wave properties are only dominant in very small particles.

## De Broglie Wavelength

Momentum of a photon - inverse to wavelength.

$$p = \frac{E}{c} \quad \text{Einstein's special relativity theory}$$

since  $E = \frac{hc}{\lambda}$  

$$p = \frac{h}{\lambda}$$

Wavelength of a particle- inverse to momentum.


$$\lambda = \frac{h}{p}$$

De Broglie proposed that this wavelength applied to material particles as well as for photons. (1924)



Lois De Boglie

## Wave properties



Diffraction of waves  
 Increases as the ratio  $\lambda/a$

## de Broglie wavelength of a baseball




$m = 0.15 \text{ kg}$   
 $v = 45 \text{ m/s}$

A baseball with a mass of 0.15 kg is pitched at 45 m/s  
 What is its De Broglie wavelength?

$$\lambda = \frac{h}{p} = \frac{h}{mv} = \frac{6.6 \times 10^{-34} \text{ Js}}{(0.15 \text{ kg})(45 \text{ m/s})} = 9.8 \times 10^{-35} \text{ m}$$

Diffraction effects of a baseball are negligible

## de Broglie wavelength of an electron



$m = 9.1 \times 10^{-31} \text{ kg}$

Find the de Broglie wavelength of a 1000 eV electron.

$$KE = \frac{1}{2}mv^2 = \frac{m^2v^2}{2m} = \frac{p^2}{2m} = eV$$

$$\lambda = \frac{h}{p} = \frac{h}{\sqrt{2meV}}$$

$V = 1000 \text{ V}$

atomic dimensions


$$\lambda = \frac{6.6 \times 10^{-34} \text{ Js}}{\sqrt{2(9.1 \times 10^{-31} \text{ kg})(1.6 \times 10^{-19} \text{ C})(1000 \text{ J/C})}} = 4.0 \times 10^{-11} \text{ m}$$

## Proof of the wave nature of electrons by Electron Diffraction

- Davisson – Germer Experiment
- Thompson Experiment

Showed the wave nature of light by diffraction of electrons by crystals.

## Diffraction of light from crystals




Crystals act as a three-dimensional diffraction grating  
Light with wavelength close to the inter-atomic spacing (x-rays) is diffracted.

## Diffraction of electrons from crystals

Davisson-Germer Experiment (1927)

An electron beam is scattered from a crystal



The scattered beam shows a diffraction pattern expected for the crystal spacing.

## Comparison between electron diffraction and x-ray diffraction


George Thomson (1927)

Aluminum powder Diffraction pattern

Either x-rays or electrons

The electron wavelength was adjusted to the same value as the x-ray by varying the voltage.


The diffraction pattern for x-rays and electrons are very similar.



X-ray pattern electron pattern

Material particles have wave properties.


## Electron microscopy




$$\lambda = \frac{h}{mv}$$

Shorter wavelength can be obtained by increasing  $v$ , the speed of the electron.

## Electron microscopy




electron microscope pictures of ribosomes



model of ribosome

Electron microscopy can be used to image structures of molecules.

## Diffraction of particles Probabilistic Interpretation of the wave amplitude.




The amplitude<sup>2</sup> is interpreted as the **probability** of the particle hitting the screen at a certain position  
This is true for electrons as well as photons.

## Wavefunction

In quantum mechanics the result of an experiment is given in terms of a wavefunction  $\Psi$ . The square of the wavefunction  $\Psi^2$  is the probability of the particle being at a certain position.

The wavefunction can be calculated using using the Schrödinger Equation. For instance for electrons in an atom.

## Wave property of particles




Decreasing the slit reduces  $\Delta x$   
But increases the width of the diffraction,  $\Delta v_x$

When  $\Delta x$  decreases,  $\Delta p_x$  increases.

## Uncertainty Principle

Particle passing through a slit --  
The uncertainty in position is  $\Delta x$

The uncertainty in the x component of momentum is  $\Delta p_x = m\Delta v_x$



The particle is diffracted

$$\Delta x \sin \theta = \lambda = \frac{h}{mv}$$

$$\sin \theta = \theta = \frac{\Delta v_x}{v}$$

$$\Delta x \frac{\Delta v_x}{v} = \frac{h}{mv}$$

Therefore  $\Delta x \Delta p_x = h$


most often written as an inequality

$$\Delta x \Delta p_x \geq \frac{h}{4\pi}$$

The position and velocity cannot be known with unlimited certainty.

## The size of an atom

What accounts for the size of the hydrogen atom?



$$PE = -\frac{k_0 e^2}{r}$$

PE  $\rightarrow$  - infinity as  $r \rightarrow$  zero

Classical electrostatics predicts that the potential energy of the hydrogen atom should go to -infinity

The finite size of the atom is a quantum mechanical effect.

As the size of the atom  $r$  decreases – The uncertainty in the momentum increases until the uncertainty in momentum limits the momentum. Then **the kinetic energy must increase** due to the uncertainty principle.

Use linear momentum as a rough estimate.

$$\Delta x \Delta p_x \approx h$$

$$\Delta p_x \approx \frac{h}{\Delta x} = \frac{h}{r}$$




$$\Delta p_x = \frac{h}{2r} \approx p_x \quad p_x \text{ cannot be smaller than } \Delta p_x$$

$$KE = \frac{1}{2} m v_x^2 = \frac{p_x^2}{2m} \approx \frac{h^2}{8r^2 m}$$


KE increases as  $1/r^2$

$E = KE + PE$  goes through a minimum as a function of  $r$




## Tunneling across a barrier

a macroscopic object impinging on a barrier the object cannot penetrate within the barrier.





a wave particle impinging on a barrier can penetrate within the barrier for distance. and go through the barrier if it is thin enough.



The probability of tunneling decreases exponentially with the width of the barrier.


## Electron Tunneling Microscope




$$\text{Electron\_Probability} \propto |\Psi|^2 \approx \Psi_0^2 e^{-2\alpha r}$$

$$\alpha \sim 10 \text{ nm}^{-1}$$

The tunneling probability falls off exponentially with  $r$  and is a sensitive function of distance



## Tunneling in Photosynthesis



Bacterial Reaction Center

Electron transfer time vs distance

Electron transfer is due to tunneling. Transfer times decay exponentially as a function of distance.