

9.1 Nuclear Physics

Properties of nuclei
 Binding Energy
 Radioactive decay
 Natural radioactivity

Properties of the nucleus

Consists of protons and neutrons

Z = no. of protons (Atomic number)
 N = no. of neutrons (Neutron number)
 A = Z+N (Mass number)

Notation :For element X with mass no. A and Atomic no. Z



Example ${}^4_2\text{He}$



Isotopes

Isotopes are nuclei that have the same no. of protons but different no. of neutrons.

The chemical properties are the same but the nuclear properties are different. i.e. some isotopes may be unstable and are radioactive.

eg. ${}^1_1\text{H}$ Hydrogen - stable
 ${}^2_1\text{H}$ Deuterium - stable
 ${}^3_1\text{H}$ Tritium - radioactive


Stable Nuclei

Plot of N vs Z for stable nuclei

Outline shows unstable nuclei

Some radioactive nuclei

${}^{14}_6\text{C}$ ${}^{32}_{15}\text{P}$ ${}^{131}_{53}\text{I}$ ${}^{238}_{92}\text{U}$



Size of the nucleus

From scattering experiments with alpha particles He^{++}



For gold nuclei

closest $r < 3 \times 10^{-14} \text{ m}$

This is smaller than the size of atoms $r \sim 10^{-10} \text{ m}$

Size of the nucleus

Radius varies as the cube root of A

$$r = r_0 A^{1/3}$$

where $r_0 = 1.2 \times 10^{-15} \text{ m}$

example

For Uranium 238, ${}^{238}_{92}\text{U}$
 A=238


$$r = 1.2 \times 10^{-15} (238)^{1/3} = 7.4 \times 10^{-15} \text{ m}$$



Forces in the nuclei

Coulomb forces

The protons repel each other with Coulomb forces. These are enormously large due to the small size.



Nuclear forces

The nucleus is held together by the nuclear force. This force acts only at **short range** ($\sim 10^{-15}$ m) and is independent of charge (i.e. acts between proton-proton, proton-neutron and neutron-neutron).

Equivalence of mass and energy

A famous result from Einstein's Special Relativity Theory

$$E = mc^2$$

mass can be converted into energy

Energy equivalent of an electron mass

$$E = mc^2 = (9.1 \times 10^{-31} \text{ kg})(3 \times 10^8 \text{ m/s})^2 = 8.2 \times 10^{-14} \text{ J} \\ = 5.1 \times 10^5 \text{ eV} = 0.51 \text{ MeV}$$

An electron can be annihilated (converted completely to energy). A 0.51 MeV photon is produced.

Mass changes when energy is lost or gained

Gasoline + $O_2 \rightarrow CO_2$ + water + kinetic energy (heat)
 $\Sigma mc^2 + \Sigma \text{Energy} = \text{constant}$

The energy released is equal to the change in mass in the reaction.

$$E = \Delta mc^2$$

CO_2 + water is lighter.

Burning 1 kg of gasoline releases 44×10^6 J of energy.



The change in mass is

$$\Delta m = \frac{E}{c^2} = \frac{44 \times 10^6 \text{ J}}{(3 \times 10^8 \text{ m/s})^2} = 5 \times 10^{-10} \text{ kg} \quad \text{small change in mass}$$

Binding energy

The binding energy of the nucleus can be determined by measuring the mass of the components and the final product.

$$E = \Delta mc^2$$



$\Delta m = \text{mass (hydrogen atom)} + \text{mass (neutron)} - \text{mass (deuterium atom)}$


$$\Delta m = 1.007825u + 1.008665u - 2.014102u = 2.39 \times 10^{-3}u \\ u = 1.660559 \times 10^{-27} \text{ kg (atomic mass unit)}$$

$$E = \Delta mc^2 = (2.39 \times 10^{-3}u)(1.66 \times 10^{-27} \text{ kg/u})(3 \times 10^8 \text{ m/s})^2 = 3.6 \times 10^{-13} \text{ J}$$

$$E = 2.2 \times 10^6 \text{ eV} = 2.2 \text{ MeV} \quad \text{Binding energy of the deuteron}$$


Binding energy per nucleon (E/A)

N neutrons + Z protons \rightarrow Atom + energy



Goes through a maxima at ${}^{56}\text{Fe}$

Binding energy per nucleon (E/A)



Goes through a maxima at ${}^{56}\text{Fe}$

Fusion (combine small nuclei) – increases binding energy

Fission (break large nuclei) – increases binding energy

Radioactivity

Unstable nuclei decay releasing energy and radiation.


Three types of radiation

alpha (α) particles - ${}^4_2\text{He}$ nuclei (+ charge)
 beta (β) particles - electrons (- charge)
 gamma (γ) particles - high frequency electromagnetic radiation. (uncharged)


Increasing penetration

Decay process

Alpha decay




Uranium decay




Beta Decay

electron e^-




positron e^+




e^+ positron has the same mass as an electron but a positive charge. the antiparticle of the electron
 γ neutrino, no charge, low mass particle – carries of some momentum and energy
 $\bar{\gamma}$ anti-neutrino, antiparticle of the neutrino

Beta decay

electron



positron



Gamma Decay

Many nuclear reaction produce nuclear excited states that decay giving of high energy electromagnetic radiation-gamma rays.



Radiation

Penetration depth


alpha particles – Stopped by a sheet of paper

beta particles - Stopped by a mm of aluminum

gamma particles - Stopped by a few cm of lead

Measuring radiation

Geiger Counter



ionization of gas causes current flow.


Natural radioactivity

Many elements found in nature are unstable and decay emitting radioactivity.

These include Uranium, ^{238}U , Radon ^{224}Ra and Potassium ^{40}K . Carbon ^{14}C ,

Natural radioactive decay

Thorium decay gives a variety of unstable products.



Cosmic rays


Cosmic rays –High energy particles (protons, alpha particles, atomic nuclei) from distant stars collide with atoms in the atmosphere and break them apart to sub atomic particles.

Some particles e.g. muons rain down to the earth's surface and are a source of background radiation.

Cloud chamber

A cloud chamber can be used to visualize radioactive particles.

The chamber contains a supersaturated vapor



The radioactive particle causes ionization of atoms. The charge initiates the condensation of the vapor, leaving a visible track.

Cloud chamber demo

<http://www.youtube.com/watch?v=kuzWNOUqLmQ>

Decay rate

The rate of decay, R, for N nuclei is proportional to N

$$R = \frac{\Delta N}{\Delta t} = \lambda N$$

λ = decay constant (units time⁻¹)

Activity – (measure of the rate of radioactive decay)

Units Curie,

$$1\text{Ci} = 3.7 \times 10^{10} \text{ Decays/s}$$

Radioactive decay

Radioactive decay is a random process the amount of material remaining varies exponentially with time.


$$N = N_0 e^{-\lambda t}$$

Decay constant λ (units 1/time)

This can also be expressed as

$$N = N_0 \left(\frac{1}{2}\right)^{t/T_{1/2}}$$

$$\text{Half Life } T_{1/2} = \frac{\ln 2}{\lambda} = \frac{0.693}{\lambda}$$



Example 29.3

The half life of radium Ra is 1.6×10^3 yr. If the sample contains 3.00×10^{16} nuclei. Find the activity in curies. (1Ci = 3.7×10^{10} decays/s)

$$T_{1/2} = \frac{\ln 2}{\lambda} = \frac{0.693}{\lambda}$$

$$\lambda = \frac{0.693}{T_{1/2}} = \frac{0.693}{1.6 \times 10^3 \text{ yr} (365 \text{ day / yr}) (24 \text{ hr / day}) (3600 \text{ s / hr})}$$

$$= 1.37 \times 10^{-11} \text{ s}^{-1}$$

$$R = \lambda N$$

$$R = 1.37 \times 10^{-11} \text{ s}^{-1} (3.00 \times 10^{16} \text{ nuclei}) = 4.12 \times 10^5 \text{ decays/s}$$

$$R = 1.1 \times 10^{-5} \text{ Ci or } 11 \mu\text{curies} \quad 3.7 \times 10^{10} \text{ decays/sCi}$$

Example 29.3

The half life of radium Ra is 1.6×10^3 yr. If the sample contains 3.00×10^{16} nuclei. Find the number of nuclei after 4.8×10^3 yr.

$$N = N_0 \left(\frac{1}{2}\right)^{t/T_{1/2}}$$

$$N = 3.00 \times 10^{16} \left(\frac{1}{2}\right)^{4.8 \times 10^3 / 1.6 \times 10^3}$$

$$N = 3.00 \times 10^{16} (1/8) = 3.75 \times 10^{15} \text{ nuclei}$$

Example 29.3

The half life of radium Ra is 1.6×10^3 yr. If the sample contains 3.00×10^{16} nuclei. Find the activity after 4.8×10^3 yr.


$$R = \lambda N$$

After this time since the no. of nuclei is reduced by a factor of 8
the decay rate will also be reduced by a factor of 8.


$$R = 11 \mu\text{Ci} / 8 = 1.4 \mu\text{Ci}$$

Radioactive dating

¹⁴C is continually formed by cosmic rays in the upper atmosphere.




so that the concentration of ¹⁴C is relatively constant over long times (longer than the half life).



The fraction of ¹⁴C decays exponentially with time.

Smoke Detector

Ionization of air by a radioactive source produces a current. Smoke traps the electrons and reduces the current, setting off the alarm.



Medical Applications.

Radiation Damage.

- Nuclear particles have much higher energies than chemical bonds.
- Radiation breaks chemical bonds – forming reactive chemical species – radicals.
- Reactive chemicals cause radiation damage to biological systems – often reaction with DNA

Radiation Therapy


Radiation is often used in treating cancer.



external radiation

Radioimmunotherapy

New methods for can deliver radiation more specifically to target cells



Treatment of non-Hodgkins lymphoma with radioimmunotherapy

Properties of ^{131}I

Iodine 131

Half-life – 8.07 days

Beta particle
 maximum energy- 807 keV
 average energy - 182 keV

Range in tissue -2.4 mm

Common clinical applications
 Radioimmunotherapy, thyroid ablation for benign and malignant disease

Medical Imaging

- X-ray Computer axial tomography (CAT)
- Positron emission tomography (PET)
- Magnetic resonance imaging (MRI)
- Contrast
- Resolution.


CAT scan

Contrast – x-ray absorption may use heavy elements to increase contrast i.e I, Ba



A three- dimensional image is reconstructed from many two dimensional pictures.

Computer tomography




at each detector absorbance is due to the sum of the absorptions from each segment.

$$A (\text{detector}) = A_1 + A_2$$

For 4 detectors -> 4 linear equations and 4 segment -> 4 unknowns \Rightarrow Solve for each absorbance


CAT scan of a chest





Positron Emission Tomography

- Emission of 2 gamma ray photons traveling in opposite direction by Positron-Electron annihilation. (conservation of momentum)
- Positrons are produced by decay of short lived radioactive nuclei such as


^{18}F ($T_{1/2}=110$ min)



Annihilation produces 2 0.51 MeV photons




PET imaging system



Two coincident detectors are used to detect the gamma rays. The source is in a line directly between the two detectors.

PET scan of the human brain



^{15}O ($T_{1/2}=2$ min) marks the consumption of O_2 due to brain activity.