

8-1 $E = \frac{\hbar^2 \pi^2}{2m} \left[\left(\frac{n_1}{L_x} \right)^2 + \left(\frac{n_2}{L_y} \right)^2 + \left(\frac{n_3}{L_z} \right)^2 \right]$

$L_x = L$, $L_y = L_z = 2L$. Let $\frac{\hbar^2 \pi^2}{8mL^2} = E_0$. Then $E = E_0(4n_1^2 + n_2^2 + n_3^2)$. Choose the quantum numbers as follows:

n_1	n_2	n_3	$\frac{E}{E_0}$		
1	1	1	6		ground state
1	2	1	9	*	first two excited states
1	1	2	9	*	
2	1	1	18		
1	2	2	12	*	next excited state
2	1	2	21		
2	2	1	21		
2	2	2	24		
1	1	3	14	*	next two excited states
1	3	1	14	*	

Therefore the first 6 states are ψ_{111} , ψ_{121} , ψ_{112} , ψ_{122} , ψ_{113} , and ψ_{131} with relative energies $\frac{E}{E_0} = 6, 9, 9, 12, 14, 14$. First and third excited states are doubly degenerate.

8-2 (a) $n_1 = 1, n_2 = 1, n_3 = 1$

$$E_0 = \frac{3\hbar^2 \pi^2}{2mL^2} = \frac{3\hbar^2}{8mL^2} = \frac{3(6.626 \times 10^{-34} \text{ Js})^2}{8(9.11 \times 10^{-31} \text{ kg})(2 \times 10^{-10} \text{ m})^2} = 4.52 \times 10^{-18} \text{ J} = 28.2 \text{ eV}$$

(b) $n_1 = 2, n_2 = 1, n_3 = 1$ or
 $n_1 = 1, n_2 = 2, n_3 = 1$ or
 $n_1 = 1, n_2 = 1, n_3 = 2$

$$E_1 = \frac{6\hbar^2}{8mL^2} = 2E_0 = 56.4 \text{ eV}$$

8-3 $n^2 = 11$

(a) $E = \left(\frac{\hbar^2 \pi^2}{2mL^2} \right) n^2 = \frac{11}{2} \left(\frac{\hbar^2 \pi^2}{mL^2} \right)$

(b)
$$\begin{array}{ccc} n_1 & n_2 & n_3 \\ \hline 1 & 1 & 3 \\ 1 & 3 & 1 & 3\text{-fold degenerate} \\ 3 & 1 & 1 \end{array}$$

(c) $\psi_{113} = A \sin\left(\frac{\pi x}{L}\right) \sin\left(\frac{\pi y}{L}\right) \sin\left(\frac{3\pi z}{L}\right)$

$$\psi_{131} = A \sin\left(\frac{\pi x}{L}\right) \sin\left(\frac{3\pi y}{L}\right) \sin\left(\frac{\pi z}{L}\right)$$

$$\psi_{311} = A \sin\left(\frac{3\pi x}{L}\right) \sin\left(\frac{\pi y}{L}\right) \sin\left(\frac{\pi z}{L}\right)$$

- 8-4 (a) $\psi(x, y) = \psi_1(x)\psi_2(y)$. In the two-dimensional case, $\psi = A(\sin k_1 x)(\sin k_2 y)$ where $k_1 = \frac{n_1\pi}{L}$ and $k_2 = \frac{n_2\pi}{L}$.

$$(b) E = \frac{\hbar^2 \pi^2 (n_1^2 + n_2^2)}{2mL^2}$$

If we let $E_0 = \frac{\hbar^2 \pi^2}{mL^2}$, then the energy levels are:

n_1	n_2	$\frac{E}{E_0}$	
1	1	1	$\rightarrow \psi_{11}$
1	2	<u>5</u> 2	$\rightarrow \psi_{12}$
2	1	<u>5</u> 2	$\rightarrow \psi_{21}$
2	2	4	$\rightarrow \psi_{22}$

- 8-5 (a) $n_1 = n_2 = n_3 = 1$ and
- $$E_{111} = \frac{3\hbar^2}{8mL^2} = \frac{3(6.63 \times 10^{-34})^2}{8(1.67 \times 10^{-27})(4 \times 10^{-28})} = 2.47 \times 10^{-13} \text{ J} \approx 1.54 \text{ MeV}$$

- (b) States 211, 121, 112 have the same energy and $E = \frac{(2^2 + 1^2 + 1^2)\hbar^2}{8mL^2} = 2E_{111} \approx 3.08 \text{ MeV}$
and states 221, 122, 212 have the energy $E = \frac{(2^2 + 2^2 + 1^2)\hbar^2}{8mL^2} = 3E_{111} \approx 4.63 \text{ MeV}$.


- (c) Both states are threefold degenerate.

8-10 $n = 4$, $l = 3$, and $m_l = 3$.


(a) $L = [l(l+1)]^{1/2} \hbar = [3(3+1)]^{1/2} \hbar = 2\sqrt{3}\hbar = 3.65 \times 10^{-34}$ Js

(b) $L_z = m_l \hbar = 3\hbar = 3.16 \times 10^{-34}$ Js

$$8-12 \quad \psi(r) = \left(\frac{1}{\pi}\right)^{1/2} \left(\frac{1}{a_0}\right)^{3/2} e^{-\frac{r}{a_0}}$$



- (b) The probability of finding the electron in a volume element dV is given by $|\psi|^2 dV$. Since the wave function has spherical symmetry, the volume element dV is identified here with the volume of a spherical shell of radius r , $dV = 4\pi r^2 dr$. The probability of finding the electron between r and $r + dr$ (that is, within the spherical shell) is $P = |\psi|^2 dV = 4\pi r^2 |\psi|^2 dr$.



$$(d) \quad \int |\psi|^2 dV = 4\pi \int |\psi|^2 r^2 dr = 4\pi \left(\frac{1}{\pi}\right) \left(\frac{1}{a_0^3}\right) \int_0^\infty e^{-2\frac{r}{a_0}} r^2 dr = \left(\frac{4}{a_0^3}\right) \int_0^\infty e^{-2\frac{r}{a_0}} r^2 dr$$

Integrating by parts, or using a table of integrals, gives

$$\int |\psi|^2 dV = \left(\frac{4}{a_0^3}\right) \left[2 \left(\frac{a_0}{2}\right)^3 \left(\frac{2}{a_0}\right)^3 \right] = 1.$$

$$(e) \quad P = 4\pi \int_n^{r_2} |\psi|^2 r^2 dr \text{ where } n = \frac{a_0}{2} \text{ and } r_2 = \frac{3a_0}{2}$$

$$\begin{aligned}
P &= \left(\frac{4}{a_0^3} \right)_{r_1}^{\infty} r^2 e^{-2\mu a_0} dr \quad \text{let } z = \frac{2r}{a_0} \\
&= \frac{1}{2} \int_1^3 z^2 e^{-z} dz \\
&= -\frac{1}{2} \left(z^2 + 2z + 2 \right) e^{-z} \Big|_1^3 \quad (\text{integrating by parts}) \\
&= -\frac{17}{2} e^{-3} + \frac{5}{2} e^{-1} = 0.496
\end{aligned}$$

8-13 $Z = 2$ for He^+

(a) For $n = 3$, l can have the values of 0, 1, 2

$$\begin{aligned}
l = 0 &\rightarrow m_l = 0 \\
l = 1 &\rightarrow m_l = -1, 0, +1 \\
l = 2 &\rightarrow m_l = -2, -1, 0, +1, +2
\end{aligned}$$

(b) All states have energy $E_3 = \frac{-Z^2}{3^2} (13.6 \text{ eV})$

$$E_3 = -6.04 \text{ eV}.$$

8-14 $Z = 3$ for Li^{2+}

(a) $n = 1 \rightarrow l = 0 \rightarrow m_l = 0$
 $n = 2 \rightarrow l = 0 \rightarrow m_l = 0$
and $l = 1 \rightarrow m_l = -1, 0, +1$

(b) For $n = 1$, $E_1 = -\left(\frac{3^2}{1^2}\right)(13.6) = -122.4 \text{ eV}$
For $n = 2$, $E_2 = -\left(\frac{3^2}{2^2}\right)(13.6) = -30.6 \text{ eV}$

8-16 For a d state, $l = 2$. Thus, m_l can take on values $-2, -1, 0, 1, 2$. Since $L_z = m_l \hbar$, L_z can be $\pm 2\hbar, \pm \hbar$, and zero.

8-17 (a) For a d state, $l = 2$

$$L = [l(l+1)]^{1/2} \hbar = (6)^{1/2} (1.055 \times 10^{-34} \text{ Js}) = 2.58 \times 10^{-34} \text{ Js}$$

(b) For an f state, $l = 3$

$$L = [l(l+1)]^{1/2} \hbar = (12)^{1/2} (1.055 \times 10^{-34} \text{ Js}) = 3.65 \times 10^{-34} \text{ Js}$$

8-18 The state is 6g

(a) $n = 6$

$$(b) \quad E_n = -\frac{13.6 \text{ eV}}{n^2} \quad E_6 = \frac{-13.6}{6^2} \text{ eV} = -0.378 \text{ eV}$$

(c) For a g -state, $l=4$

$$L = [l(l+1)]^{1/2} \hbar = (4 \times 5)^{1/2} \hbar = \sqrt{20} \hbar = 4.47 \hbar$$

(d) m_l can be $-4, -3, -2, -1, 0, 1, 2, 3$, or 4

$L_z = m_l \hbar$; $\cos \theta = \frac{L_z}{L} = \frac{m_l}{[l(l+1)]^{1/2}} \hbar = \frac{m_l}{\sqrt{20}}$
m_l -4 -3 -2 -1 0 1 2 3 4
L_z $-4\hbar$ $-3\hbar$ $-2\hbar$ $-\hbar$ 0 \hbar $2\hbar$ $3\hbar$ $4\hbar$
θ 153.4° 132.1° 116.6° 102.9° 90° 77.1° 63.4° 47.9° 26.6°

8-21 (a) $\psi_{2s}(r) = \frac{1}{4(2\pi)^{1/2}} \left(\frac{1}{a_0}\right)^{3/2} \left(2 - \frac{r}{a_0}\right) e^{-r/a_0}$. At $r = a_0 = 0.529 \times 10^{-10} \text{ m}$ we find

$$\begin{aligned} \psi_{2s}(a_0) &= \frac{1}{4(2\pi)^{1/2}} \left(\frac{1}{a_0}\right)^{3/2} (2-1)e^{-1/2} = (0.380) \left(\frac{1}{a_0}\right)^{3/2} \\ &= (0.380) \left[\frac{1}{0.529 \times 10^{-10} \text{ m}}\right]^{3/2} = 9.88 \times 10^{14} \text{ m}^{-3/2} \end{aligned}$$

(b) $|\psi_{2s}(a_0)|^2 = (9.88 \times 10^{14} \text{ m}^{-3/2})^2 = 9.75 \times 10^{29} \text{ m}^{-3}$

(c) Using the result to part (b), we get $P_{2s}(a_0) = 4\pi a_0^2 |\psi_{2s}(a_0)|^2 = 3.43 \times 10^{10} \text{ m}^{-1}$.

8-22 $R_{2p}(r) = A r e^{-r/a_0}$ where $A = \frac{1}{2(6)^{1/2} a_0^{5/2}}$

$$\begin{aligned} P(r) &= r^2 R_{2p}^2(r) = A^2 r^4 e^{-2r/a_0} \\ \langle r \rangle &= \int_0^\infty r P(r) dr = A^2 \int_0^\infty r^5 e^{-2r/a_0} dr = A^2 a_0^6 5! = 5a_0 = 2.645 \text{ \AA} \end{aligned}$$