

7-9. (a) For $n = 3$, $\ell = 0, 1, 2$

(b) For $\ell = 0, m = 0$. For $\ell = 1, m = -1, 0, +1$. For $\ell = 2, m = -2, -1, 0, +1, +2$.

(c) There are nine different m -states, each with two spin states, for a total of 18 states for $n = 3$.

7-10. (a) For $\ell = 4$

$$L = \sqrt{\ell(\ell+1)}\hbar = \sqrt{4(5)}\hbar = \sqrt{20}\hbar$$

$$m_\ell = 4\hbar$$

$$\theta_{\min} = \cos^{-1} \frac{4}{\sqrt{20}} \rightarrow \theta_{\min} = 26.6^\circ$$


(b) For $\ell = 2$

$$L = \sqrt{6}\hbar \quad m_\ell = 2\hbar$$

$$\theta_{\min} = \cos^{-1} \frac{2}{\sqrt{6}} \rightarrow \theta_{\min} = 35.3^\circ$$


7-12. (a)

$$\ell = 1$$
$$|\mathbf{L}| = \sqrt{2}\hbar$$




(b)

$$\ell = 2$$
$$|\mathbf{L}| = \sqrt{6}h$$



(c)

$$\ell = 4$$
$$|\mathbf{L}| = \sqrt{20}h$$



(d) $|\mathbf{L}| = \sqrt{\ell(\ell+1)}h$ (See diagrams above.)

7-13. $L^2 = L_x^2 + L_y^2 + L_z^2 \rightarrow L_x^2 + L_y^2 = L^2 - L_z^2 = \ell(\ell+1)h^2 - (mh)^2 = (6-m^2)h^2$

$$(a) (L_x^2 + L_y^2)_{\min} = (6 - 2^2)\hbar^2 = 2\hbar^2$$

$$(b) (L_x^2 + L_y^2)_{\max} = (6 - 0^2)\hbar^2 = 6\hbar^2$$

$$(c) L_x^2 + L_y^2 = (6 - 1)\hbar^2 = 5\hbar^2 \quad L_x \text{ and } L_y \text{ cannot be determined separately.}$$

$$(d) n = 3$$

$$7-15. \quad \mathbf{L} = \mathbf{r} \times \mathbf{p} \quad \frac{d\mathbf{L}}{dt} = \frac{d\mathbf{r}}{dt} \times \mathbf{p} + \mathbf{r} \times \frac{d\mathbf{p}}{dt}$$

$$\frac{d\mathbf{r}}{dt} \times \mathbf{p} = \mathbf{v} \times m\mathbf{v} = m\mathbf{v} \times \mathbf{v} = 0 \quad \text{and} \quad \mathbf{r} \times \frac{d\mathbf{p}}{dt} = \mathbf{r} \times \mathbf{F}. \quad \text{Since for } V = V(r), \text{ i.e., central forces,}$$

$$\mathbf{F} \text{ is parallel to } \mathbf{r}, \text{ then } \mathbf{r} \times \mathbf{F} = 0 \quad \text{and} \quad \frac{d\mathbf{L}}{dt} = 0$$

$$7-16. (a) \text{ For } \ell = 3, n = 4, 5, 6, \dots \text{ and } m = -3, -2, -1, 0, 1, 2, 3$$

$$(b) \text{ For } \ell = 4, n = 5, 6, 7, \dots \text{ and } m = -4, -3, -2, -1, 0, 1, 2, 3, 4$$

$$(c) \text{ For } \ell = 0, n = 1 \text{ and } m = 0$$

$$(d) \text{ The energy depends only on } n. \text{ The minimum in each case is:}$$

$$E_4 = -13.6eV/n^2 = -13.6eV/4^2 = -0.85eV$$

$$E_5 = -13.6eV/5^2 = -0.54eV$$

$$E_1 = -13.6eV$$

$$7-17. (a) 6f \text{ state: } n = 6, \ell = 3$$

$$(b) E_6 = -13.6eV/n^2 = -13.6eV/6^2 = -0.38eV$$

$$(c) L = \sqrt{\ell(\ell+1)}\hbar = \sqrt{3(3+1)}\hbar = \sqrt{12}\hbar = 3.65 \times 10^{-34} \text{ J}\cdot\text{s}$$

$$(d) L_z = m\hbar \quad L_z = -3\hbar, -2\hbar, -1\hbar, 0, 1\hbar, 2\hbar, 3\hbar$$

7-21. (a) For the ground state, $P(r)\Delta r = \psi^2(4\pi r^2)\Delta r = \frac{4r^2}{a_0^3}e^{-2r/a_0}\Delta r$

For $\Delta r = 0.03a_0$, at $r = a_0$ we have $P(r)\Delta r = \frac{4a_0^2}{a_0^3}e^{-2}(0.03a_0) = 0.0162$

(b) For $\Delta r = 0.03a_0$, at $r = 2a_0$ we have $P(r)\Delta r = \frac{4(2a_0)^2}{a_0^3}e^{-4}(0.03a_0) = 0.0088$

7-22. $P(r) = Cr^2e^{-2Zr/a_0}$ For $P(r)$ to be a maximum,

$$\frac{dP}{dr} = C \left[r^2 \left(-\frac{2Z}{a_0} \right) e^{-2Zr/a_0} + 2r e^{-2Zr/a_0} \right] = 0 \rightarrow C \times \frac{2Zr}{a_0} \left(\frac{a_0}{Z} - r \right) e^{-2Zr/a_0} = 0$$

This condition is satisfied with $r = 0$ or $r = a_0/Z$. For $r = 0$, $P(r) = 0$ so the maximum $P(r)$ occurs for $r = a_0/Z$.

7-27. For the most likely value of r , $P(r)$ is a maximum, which requires that (see Problem 7-25)

$$\frac{dP}{dr} = A \cos^2 \theta \left[r^4 \left(-\frac{Z}{a_0} \right) e^{-Zr/a_0} + 4r^3 e^{-Zr/a_0} \right] = 0$$

For hydrogen $Z = 1$ and $A \cos^2 \theta (r^3/a_0)(4a_0 - r)e^{-r/a_0} = 0$. This is satisfied for $r = 0$ and $r = 4a_0$. For $r = 0$, $P(r) = 0$ so the maximum $P(r)$ occurs for $r = 4a_0$.

7-65. $\psi_{100} = \frac{1}{\sqrt{\pi}} \left(\frac{Z}{a_0} \right)^{3/2} e^{-Zr/a_0}$ (Equations 7-30 and 7-31)

$$P(r) = 4\pi r^2 \psi_{100}^* \psi_{100} \quad (\text{Equation 7-32})$$

$$= 4\pi r^2 \frac{Z^3}{\pi a_0^3} e^{-2Zr/a_0} = \frac{4Z^3}{a_0^3} r^2 e^{-2Zr/a_0}$$

$$\langle r \rangle = \int_0^\infty r P(r) dr = \int_0^\infty \frac{4Z^3}{a_0^3} r^3 e^{-2Zr/a_0} dr$$

$$= \frac{a_0}{4Z} \int_0^\infty \left(\frac{2Zr}{a_0} \right)^3 e^{-2Zr/a_0} d(2Zr/a_0) = \frac{a_0}{4Z} \times 3! = \frac{3a_0}{2Z}$$

7-68. $\theta_{\min} = \cos^{-1} \left[m_\ell \hbar / \sqrt{\ell(\ell+1)} \hbar \right]$ with $m_\ell = \ell$.

$\cos \theta_{\min} = \ell / \sqrt{\ell(\ell+1)}$. Thus, $\cos^2 \theta_{\min} = \ell^2 / [\ell(\ell+1)] = 1 - \sin^2 \theta_{\min}$

or, $\sin^2 \theta_{\min} = 1 - \frac{\ell^2}{\ell(\ell+1)} = \frac{\ell(\ell+1) - \ell^2}{\ell(\ell+1)} = \frac{\ell^2 + \ell - \ell^2}{\ell(\ell+1)}$

And, $\sin \theta_{\min} = \left(\frac{1}{\ell+1} \right)^{1/2}$ For large ℓ , θ_{\min} is small.

Then $\sin \theta_{\min} \approx \theta_{\min} = \left(\frac{1}{\ell+1} \right)^{1/2} \approx \frac{1}{(\ell)^{1/2}}$

7-70. $P(r) = \frac{4Z^3}{a_0^3} r^2 e^{-2Zr/a_0}$ (See Problem 7-65)

For hydrogen, $Z = 1$ and at the edge of the proton $r = R_0 = 10^{-15} m$. At that point, the exponential factor in $P(r)$ has decreased to:

$$e^{-2R_0/a_0} = e^{-2(10^{-15})/(0.529 \times 10^{-10} m)} = e^{-(3.78 \times 10^{-5})} \approx 1 - 3.78 \times 10^{-5} \approx 1$$

Thus, the probability of the electron in the hydrogen ground state being inside the nucleus, to better than four figures, is:

$$\begin{aligned} P(r) &= \frac{4r^2}{a_0^3} & P &= \int_0^{R_0} P(r) dr = \int_0^{R_0} \frac{4r^2}{a_0^3} = \frac{4}{a_0^3} \int_0^{R_0} r^2 dr = \frac{4}{a_0^3} \left. \frac{r^3}{3} \right|_0^{R_0} \\ & & &= \frac{4}{a_0^3} \left(\frac{R_0^3}{3} \right) = \frac{4(10^{-15} m)^3}{3(0.529 \times 10^{-10} m)^3} = 9.0 \times 10^{-15} \end{aligned}$$

7-72. (a) Substituting $\psi(r, \theta)$ into Equation 7-9 and carrying out the indicated operations yields (eventually)

$$-\frac{\hbar^2}{2\mu} \psi(r, \theta) \left[2/r^2 - 1/4a_0^2 \right] - \frac{\hbar^2}{2\mu} \psi(r, \theta) (-2/r^2) + V\psi(r, \theta) = E\psi(r, \theta)$$

Canceling $\psi(r, \theta)$ and recalling that $r^2 = 4a_0^2$ (because ψ given is for $n = 2$) we

$$\text{have } -\frac{\hbar^2}{2\mu}(-1/4a_0^2) + v = E$$

The circumference of the $n = 2$ orbit is: $C = 2\pi(4a_0) = 2\lambda \rightarrow a_0 = \lambda/4\pi = 1/2k$.

$$\text{Thus, } -\frac{\hbar^2}{2\mu}\left(-\frac{1}{4/4k^2}\right) + V = E \rightarrow \frac{\hbar^2 k^2}{2\mu} + V = E$$

(b) or $\frac{p^2}{2m} + v = E$ and Equation 7-9 is satisfied.

$$\int_0^\infty \psi^2 dx = \int A^2 \left(\frac{r}{a_0}\right)^2 e^{-r/a_0} \cos^2 \theta r^2 \sin \theta dr d\theta d\phi = 1$$

$$A^2 \int_0^\infty \left(\frac{r}{a_0}\right)^2 e^{-r/a_0} r^2 dr \int_0^\pi \cos^2 \theta \sin \theta d\theta \int_0^{2\pi} d\phi = 1$$

Integrating (see Problem 7-23),

$$A^2 (6a_0^3)(2/3)(2\pi) = 1$$

$$A^2 = 1/8a_0^3\pi \rightarrow A = \sqrt{1/8a_0^3\pi}$$