

3-60. Let $x = \frac{\varepsilon}{kT} = \frac{hf}{kT}$ in Equation 3-15:

$$\sum_{n=0}^{\infty} f_n = A \sum_{n=0}^{\infty} e^{-nx} = A \left[e^0 + e^{-x} + (e^{-x})^2 + (e^{-x})^3 + \dots \right] = A(1 + y + y^2 + y^3 + \dots) = 1$$

Where $y = e^{-x}$. This sum is the series expansion of

$$(1 - y)^{-1}, \text{ i.e., } (1 - y)^{-1} = 1 + y + y^2 + y^3 + \dots. \text{ Then } \sum f_n = A(1 - y)^{-1} = 1 \text{ gives } A = 1 - y.$$

Writing Equation 3-16 in terms of x and y .

$$\bar{E} = \sum_{n=0}^{\infty} E_n A e^{-E_n/kT} = A \sum_{n=0}^{\infty} nhf e^{-nhf/kT} = Ahf \sum_{n=0}^{\infty} ne^{-nx}$$

Note that $\sum ne^{-nx} = -(d/dx) \sum e^{-nx}$. But $\sum e^{-nx} = (1 - y)^{-1}$, so we have

$$\sum ne^{-nx} = -\frac{d}{dx} \sum e^{-nx} = -\frac{d}{dx} (1 - y)^{-1} = (1 - y)^{-2} \left(-\frac{dy}{dx} \right) = y(1 - y)^{-2}$$

$$\text{Since } \frac{dy}{dx} = \frac{d(e^{-x})}{dx} = -e^{-x} = -y.$$

Multiplying this sum by hf and by $A = (1 - y)$, the average energy is

$$\bar{E} = hfA \sum_{n=0}^{\infty} ne^{-nx} = hf(1 - y)y(1 - y)^{-2} = \frac{hfy}{1 - y} = \frac{hfe^{-x}}{1 - e^{-x}}$$

Multiplying the numerator and the denominator by e^{-x} and substituting for x , we obtain

$$\bar{E} = \frac{hf}{e^{hf/kT} - 1}, \text{ which is Equation 3-17.}$$