

Toric and their Destructors

- integrability \Rightarrow can write as action-angle form:

$$\begin{cases} \frac{d\underline{I}}{dt} = \underline{\omega} \cdot \frac{d\underline{\theta}}{dt} = \underline{\omega}(\underline{I}) \\ \text{const } \underline{I}. \end{cases}$$

\Rightarrow motion defines toric



$$\frac{d\underline{\theta}}{dt} = \underline{\omega}_1(\underline{I}_1) t$$

$$\frac{d\underline{\phi}}{dt} = \underline{\omega}_2(\underline{I}_2) t$$

scanning $\underline{I}_1, \underline{I}_2$ (linked to E)
 \Rightarrow

define nested toric



etc. $\left\{ \begin{array}{l} \text{eg. box} \\ \omega_1 = \pi^2 \underline{I}_1 / m a^2 \\ \omega_2 = \pi^2 \underline{I}_2 / m b^2 \end{array} \right.$

- motion on each toroidal surface will cover surface ergodically, unless $\underline{\omega}_1$ rational.
 - many surfaces \Rightarrow define volume of phase space,
- $$E = \underline{I}_1 \omega_1 + \underline{I}_2 \omega_2$$

- motion is conditionally periodic

i.e. ergodic motion on bounded surface
 \Rightarrow Poincaré recurrence guarantees nearby return to i.c.

\Rightarrow How robust are toroidal surfaces?

i.e. if $H \rightarrow H_0(\underline{I}) + \epsilon H(\underline{I}, \underline{\phi})$

\uparrow
 symmetry breaking
 perturbation

can we integrate the perturbed system to some order in ϵ ?

i.e. transform $\underline{I}, \underline{\phi} \rightarrow \underline{J}, \underline{\phi}$

s/t $\left. \begin{array}{l} \dot{\underline{J}} = 0 \\ \dot{\underline{\phi}} = \omega(\underline{J}) \end{array} \right\}$ to specified order in P.T.?

This is equivalent to exploring "fragility of surfaces" \Rightarrow i.e. can nested structure be maintained with $o(\epsilon)$ deformation?

n.b. \rightarrow intro to canonical perturbation theory

→ start with 7 deg freedom!

$$J = I + o(\epsilon)$$

$$q = \phi + o(\epsilon)$$

then: old: I, ϕ

new: J, ϕ

oft $j = 0$
to $o(\epsilon)$

so have C-T. problem:

$$p \leftrightarrow I$$

$$q \leftrightarrow \phi$$

(old)

$$p = J$$

$$q = \phi$$

(new)

so

index

$$q \leftrightarrow \phi$$

$$p \leftrightarrow J$$

def

$$p \leftrightarrow I$$

$$q = \phi$$

$$p = \frac{\partial F}{\partial q} = \frac{\partial S}{\partial q}$$

$$F = S$$

here,

$$S = H - J$$

fctn.

so

$$I = \partial S / \partial \theta$$

$$\phi = \partial S / \partial J$$

where: $S = S_0 + \epsilon S_1$ ↗ unknown

$$= J\theta + \epsilon S_1$$

now here:

$$S = S_0 + \epsilon S_1$$

$$H'(J) \equiv K(J)$$

new, integrated Hamiltonian \rightarrow fctn of J , only

↗ no-label.

and can expand:

$$K(J) = K_0(J) + \epsilon K_1(J) + \dots$$

∞

$$K(J) = H(I, \theta)$$

$$= H_0\left(\frac{\partial S}{\partial \theta}, \theta\right) + \epsilon H_1\left(\frac{\partial S}{\partial \theta}, \theta\right) + \dots$$

n.b:
$$\begin{aligned} \mathcal{S}' &= \mathcal{S}_0 + \epsilon \mathcal{S}_1 \\ &= \mathcal{J}\mathcal{Q} + \epsilon \mathcal{S}_1 \end{aligned}$$

$$I = \mathcal{J} + \epsilon \frac{\partial \mathcal{S}_1}{\partial \mathcal{Q}} \quad \Rightarrow \quad \mathcal{J} = I - \epsilon \frac{\partial \mathcal{S}_1}{\partial \mathcal{Q}}$$

$$\phi = \mathcal{Q} + \epsilon \frac{\partial \mathcal{S}_1}{\partial \mathcal{J}} \quad \phi = \mathcal{Q} + \epsilon \frac{\partial \mathcal{S}_1}{\partial \mathcal{J}}$$

now, plugging \mathcal{J} in to relation for H' etc

$$K_0(\mathcal{J}) + \epsilon K_1(\mathcal{J}) + \epsilon^2 K_2(\mathcal{J})$$

$$= H_0 \left(\mathcal{J} + \epsilon \frac{\partial \mathcal{S}_1}{\partial \mathcal{Q}} + \epsilon^2 \frac{\partial \mathcal{S}_2}{\partial \mathcal{Q}} + \dots \right)$$

$$+ \epsilon H_1 \left(\mathcal{J}_1 + \epsilon \frac{\partial \mathcal{S}_1}{\partial \mathcal{Q}} + \dots, \mathcal{Q} \right)$$

cranking expansion to $O(\epsilon^2)$:

$$K_0(J) + \epsilon K_1(J) + \epsilon^2 K_2(J) + \dots =$$

$$H_0(J) + \epsilon \frac{\partial S_1}{\partial \theta} \frac{\partial H_0}{\partial J} + \epsilon^2 \frac{\partial S_2}{\partial \theta} \frac{\partial H_0}{\partial J}$$

$$+ \epsilon H_1(J, \theta) + \epsilon^2 \frac{\partial S_1}{\partial \theta} \frac{\partial H_1(J)}{\partial J}$$

$$+ \frac{1}{2} \epsilon^2 \left(\frac{\partial S_1}{\partial \theta} \right)^2 \frac{\partial^2 H_0}{\partial J^2}$$

matching order-by-order:

$$H_0 = H_0$$

$$K_1(J) = \frac{\partial S_1}{\partial \theta} \frac{\partial H_0}{\partial J} + H_1(J, \theta)$$

$$K_2(J) = \frac{1}{2} \left(\frac{\partial S_1}{\partial \theta} \right)^2 \frac{\partial^2 H_0}{\partial J^2} + \frac{\partial S_2}{\partial \theta} \frac{\partial H_0}{\partial J}$$

$$+ \frac{\partial S_1}{\partial \theta} \frac{\partial H_1}{\partial J} + H_2$$

etc.

if θ present.

For $\theta(\epsilon)$:

$$\begin{aligned}
 K_1(J) &= \frac{\partial S_1}{\partial \theta} \frac{\partial H_0}{\partial J} + H_1(J, \theta) \\
 &= \frac{\partial S_1}{\partial \theta} \omega_0(J) + H_1(J, \theta)
 \end{aligned}$$

\downarrow
 winding frequency

where understand:

$$\begin{aligned}
 I &= J + \theta(\epsilon) \\
 \phi &= \theta + \theta(\epsilon)
 \end{aligned}$$

$$\begin{aligned}
 \theta &= \phi - \epsilon \frac{\partial S_1}{\partial J} \\
 \mathbf{I} &= J + \epsilon \frac{\partial S_1}{\partial \theta}
 \end{aligned}$$

Now, if define:

$$H_1 = \underbrace{\langle H_1 \rangle}_{\text{avg.}} + \underbrace{\tilde{H}_1}_{\substack{\text{dep piece} \\ \text{(symmetry breaking)}}}$$

$$\langle H_1 \rangle = \int_0^{2\pi} \frac{d\theta}{2\pi} H_1$$

(mean part)

then

averaging $K_1(J)$ eqn \Rightarrow

$$\boxed{K_1(J) = \langle H_1 \rangle}$$

and for S_1 , from solvability:

$$\begin{aligned} \omega_0(J) \frac{\partial S_1}{\partial \theta} &= K_1(J) - H_1 \\ &= \underbrace{K_1(J)}_{\sim} - \underbrace{\langle H_1 \rangle}_{\sim} - \tilde{H}_1 \\ &= -\tilde{H}_1 \end{aligned}$$

$$\boxed{\omega_0(J) \frac{\partial S_1}{\partial \theta} = -\tilde{H}_1}$$

Now, from before, as motion closed and periodic:

$$\tilde{H}_1 = \sum_{n=1}^{\infty} H_n(J) e^{in\theta}$$

$$S_1 = \sum_{n=1}^{\infty} S_n e^{in\theta}$$

$$J = J_0 + \epsilon S_1$$

\Rightarrow

$$\mathcal{J}_1 = - \sum_n \frac{H_n(\mathcal{J})}{in \omega_0(\mathcal{J})} e^{in\theta}$$

so can finally write full solution to $\mathcal{O}(\epsilon)$:

$$\begin{aligned} \phi &= \theta + \epsilon \frac{\partial \mathcal{J}_1}{\partial \mathcal{J}}(\mathcal{J}, \theta) \\ \mathcal{J} &= \mathcal{I} - \epsilon \frac{\partial \mathcal{J}_1}{\partial \theta}(\mathcal{J}, \theta) \\ \omega &= \omega_0(\mathcal{J}) + \epsilon \frac{\partial}{\partial \mathcal{J}} k_1(\mathcal{J}) \end{aligned}$$

where:

$$\begin{aligned} k_1 &= \langle H_1 \rangle \\ \mathcal{J}_1 &= \sum_n \frac{i H_n(\mathcal{J})}{n \omega_0(\mathcal{J})} e^{in\theta} \end{aligned}$$

so on 1 d.o.f; can define strategy of perturbative 'integration'.

BUT, if # d.o.f's > 1 :

$\Theta \rightarrow \underline{\Theta}$ (i.e. Θ, ϕ toroidal angles)

$$n \underline{\omega}_0(\underline{J}) \rightarrow \underline{\Lambda} \cdot \underline{\omega}_0(\underline{J})$$

$$\left(\begin{array}{l} \text{i.e. } \underline{\Lambda} \cdot \underline{\omega}_0 = n \omega_1(J_1) + m \omega_2(J_2) \\ \text{where } E = J_1 \omega_1 + J_2 \omega_2 \end{array} \right)$$

then if

$$\underline{\Lambda} \cdot \underline{\omega}_0(\underline{J}) \rightarrow 0$$

denominator
vanishes and
perturbation theory
fails

\Rightarrow welcome to
the "problem of
small divisors"

\Rightarrow identifies resonant surfaces

i.e. special surfaces of nested torus

where pitch of perturbation
 $n/m = \text{pitch of winding } \frac{\omega_2}{\omega_1}$

These seem (and are) most fragile
surfaces \downarrow

These surfaces are "resonant surfaces"

Classic example:

- tokamak field lines

$$m = n z(r)$$

$$z(r) = m/n$$

pitch of lines
↓

(note shear)

pitch of perturbation
↓

→ wave particle

$$v = \omega/k$$

n.b. here time makes resonance

particle velocity
↓

wave phase velocity
↓

$$\partial \psi / \partial t = H - H'$$


①

⇒ in vicinity of resonant surfaces, perturbative integration fails


② since actions are set ^{of measure} $\mu \rightarrow 0$ on whole #'s, resonant surfaces are in some sense special!

→ sneak preview

distortions called "islands" form
(const. H surface)



→
+ resonant
perturbation
 $m = 4$
 $n = 2$



Filamentation occurs.

$\omega_H \sim \sqrt{\delta B}$

- upshot:
- trajectory undertakes excursion from surface but remains near
 - phase space structure resembles that of pendulum.

→ caveat: secular ^{canonical} perturbation theory works for 1 resonance, only.

strategy:

- remove resonance by transformation to frame co-rotating with resonant variables
- Akin removal by frame change.
- n.b. really avg. over fast variable

- limitation to removal of 1
fast variable
i.e. works as resonance \leftrightarrow slow

Now,

$$H = H_0(\underline{I}) + \epsilon H_1(\underline{I}, \underline{\theta})$$

if resonance: $r\omega_1 - s\omega_2 \approx 0$
 \rightarrow resonance

\Rightarrow

$$\omega_1 = \frac{d\theta_1}{dt}$$

$$0 = r\theta_1 - s\theta_2 \text{ "slow"}$$

$$\omega_2 = \frac{d\theta_2}{dt}$$

so

$$\begin{aligned} (\underline{\omega} \cdot \underline{\nabla}_{\underline{\theta}}) f(\underline{\theta}) &= (\omega_1 \partial_{\theta_1} + \omega_2 \partial_{\theta_2}) f \\ &= (r\omega_1 - s\omega_2) F_{\text{is}} \end{aligned}$$

$\rightarrow \theta$, near resonance.

F dependence on θ is h.o. \rightarrow slow.

thus, before:

$$\underline{I}, \underline{\theta} \rightarrow \underline{J}, \underline{\phi}$$

now:

$$\left. \begin{array}{l} I_1, \theta_1 \\ I_2, \theta_2 \end{array} \right\} \rightarrow \left\{ \begin{array}{l} \overset{\text{slow}}{\downarrow} r \theta_1 - s \theta_2, \hat{J}_1 \\ \theta_2, \hat{J}_2 \end{array} \right.$$

2 fast \rightarrow 1 slow, 1 fast

$F = S'$ (old positions, new momenta)

$$= S(\theta_1, \theta_2; \hat{J}_1, \hat{J}_2)$$

and type 2, so:

$$S = \underbrace{(r\theta_1 - s\theta_2)}_{\text{slow}} \hat{J}_1 + \theta_2 \hat{J}_2 + \epsilon S_1$$

$$I_1 = \partial S / \partial \theta_1 = r \hat{J}_1 + \epsilon \partial S_1 / \partial \theta_1$$

$$I_2 = \partial S / \partial \theta_2 = (\hat{J}_2 - s \hat{J}_1) + \epsilon \partial S_1 / \partial \theta_2$$

$$\phi_1 = \partial S / \partial \hat{J}_1 = r \theta_1 - s \theta_2 + \epsilon \partial S_1 / \partial \hat{J}_1$$

$$\phi_2 = \partial S / \partial \hat{J}_2 = \theta_2 + \epsilon \partial S_1 / \partial \hat{J}_2$$

$$H = H_0(\underline{I}) + \epsilon H_1(\underline{I}, \underline{\theta})$$

$$H_1 = \sum_{\ell, m} H_{\ell, m}(\underline{I}) e^{i(\ell \theta_1 + m \theta_2)} \quad \ell, m \neq 0$$

but know:

$$\phi_1 = r \theta_1 - s \theta_2 + \mathcal{O}(\epsilon) \quad \text{Slow}$$

$$\phi_2 = \theta_2 + \mathcal{O}(\epsilon) \quad \text{Fast}$$

\Rightarrow

$$\theta_1 \cong (\phi_1 + s \phi_2) / r$$

$$\theta_2 \cong \phi_2$$

re-writing:

$$H_1 = \sum_{l,m} H_{l,m}(\underline{\hat{J}}) \exp \left[i \left(\frac{l}{r} \phi_1 + \frac{(ls+mr)}{r} \phi_2 \right) \right]$$

$\phi_2 \rightarrow$ fast

$\phi_1 \rightarrow$ slow

} distinction only possible
near resonance where
 $r\omega_1 = s\omega_2 \rightarrow 0$

[Now, average out fast ϕ_2
dependence, and focus on
evolution near resonance. \Rightarrow isolates
region
near resonance

Thus, will have
slow

$$K_1 = K_1(\underline{\hat{J}}, \phi_1) = \langle H_1 \rangle_{\phi_2}$$

$$\langle H_1 \rangle_{\phi_2} = \left\langle \sum_{l,m} H_{l,m}(\underline{\hat{J}}) \exp \left[i \left(\frac{l}{r} \phi_1 + \frac{(ls+mr)}{r} \phi_2 \right) \right] \right\rangle_{\phi_2}$$

on

Simply put:

$$\frac{p}{m} = \frac{-v}{s}$$

⇒

mode # pitch of
perturbation must
match pitch of
resonance

so

$$\sum_{l,m} \rightarrow \sum_{p(-v/s)}$$

⇒ sum over
all harmonics
of perturbation
resonant

∴

$$\sum_{l,m} \rightarrow \sum_p F_{-p, p, p, s}$$

upon ϕ_2 integration: $l_s = -mr$

$$\frac{p}{m} = \frac{-r}{s} \quad \text{but } r\omega_1 - s\omega_2 \sim 0$$

$\sim \frac{\omega_2}{\omega_1} \Rightarrow \frac{p}{m}$ ratio set by resonance.

so $H_1, l_0, m \rightarrow H_1, -m, r, m \quad p = \frac{-r}{s} m$

$\rightarrow H_1, -mr, ms$
relabel

$\rightarrow H_1, -pr, ps$

also $\frac{p}{s} = -\frac{m}{s}$ relabel: $-\frac{m}{s} \rightarrow -m$
 $-m \rightarrow -p$

so $\langle \rangle_{\phi_2}$ perturbation is

just harmonics of resonant pair $-r, s$.

$$\langle H_1 \rangle_{\phi_2} = \sum_{p=0}^{\infty} H_{-rp, sp} e^{-c p \phi_1}$$

18

$$\langle H \rangle = H_0(J) + \epsilon \sum_{p=0}^{\infty} H_{-rp, sp}^{(1)} e^{-c p \phi_1}$$

From C-T rules:

$$\frac{\partial \langle H \rangle}{\partial \phi_2} = 0 \Rightarrow \frac{d \vec{J}_2}{dt} = 0 \rightarrow \text{adiabatic invariant}$$

and from C-T rules:

$$I_1 = r \vec{J}_1$$

$$I_2 = \vec{J}_2 - s \vec{J}_1$$

$$\Rightarrow \vec{J}_2 = I_2 + \frac{s}{r} I_1$$

is adiabatic inv. of avgd Hamiltonian

$\phi, \dot{\phi} \rightarrow \text{res.}$

19.

so
$$\frac{d\vec{J}_2}{dt} = 0 \Rightarrow \frac{d\phi_2}{dt} = \frac{\partial \langle H \rangle}{\partial \vec{J}_2} \equiv \omega(\vec{J}_2)$$

Now, $\langle H \rangle = \langle H(\vec{J}_1, \phi_1, \vec{J}_2) \rangle$

→ For solution, need understand motion in \vec{J}_1, ϕ_1

→ without loss of generality, simplify by:

$\rho = 0, \pm 1$ harmonics only, contribute

so
$$\langle H \rangle = H_0(\vec{J}) + \epsilon H_{0,0}(\vec{J}) + 2\epsilon H_{\rho, \rho}(\vec{J}) \cos \phi$$

$$H_{-\rho, \rho} = H_{\rho, -\rho}$$

and seek motion near fixed points, as characterization.

so,
$$\begin{aligned} \dot{\vec{J}}_1 &= 0 \\ \dot{\phi}_1 &= 0 \end{aligned} \Rightarrow \text{f.p.} \Leftrightarrow \begin{aligned} \partial \langle H \rangle / \partial \phi_1 &= 0 \\ \partial \langle H \rangle / \partial \vec{J}_1 &= 0 \end{aligned}$$

these define: $\vec{J}_1 = 0$
 $\phi_1 = 0$ } fixed pts
of motion

18

$$\frac{\partial \langle H \rangle}{\partial \phi} = 0 \Rightarrow -2\epsilon H_{0,5}^{(1)} \sin \phi_1 = 0$$

$$\phi_1 = 0, \pm \pi$$

fixed pts.

and

$$\frac{\partial \langle H \rangle}{\partial \vec{J}_1} = 0 \Rightarrow \frac{\partial H_0(\vec{J}_1)}{\partial \vec{J}_1} + \epsilon \frac{\partial H_{0,0}(\vec{J}_1^2)}{\partial \vec{J}_1} + 2\epsilon \frac{\partial H_{0,5}^{(1)}}{\partial \vec{J}_1} \cos \phi_1 = 0$$

Now

$$\frac{\partial}{\partial \vec{J}_1} = \frac{\partial I_1}{\partial \vec{J}_1} \frac{\partial}{\partial I_1} + \frac{\partial I_2}{\partial \vec{J}_1} \frac{\partial}{\partial I_2}$$

C-T
rules

$$= r \frac{\partial}{\partial I_1} - s \frac{\partial}{\partial I_2}$$

so, $\partial \langle H \rangle / \partial \vec{J}_1 = 0 \Rightarrow$ re-express

$$0 = \begin{pmatrix} r \frac{\partial}{\partial I_1} & -s \frac{\partial}{\partial I_2} \end{pmatrix} H_0(\vec{I}) + \epsilon \frac{\partial}{\partial \vec{J}_1} H_{\epsilon,0} + 2\epsilon \frac{\partial H_{\epsilon,1}^{(4)}}{\partial \vec{J}_1} \cos \phi_1$$

$$= (r\omega_1 - s\omega_2) + \epsilon \left(\frac{\partial H_{\epsilon,0}}{\partial \vec{J}_1} + 2 \frac{\partial H_{\epsilon,1}^{(4)}}{\partial \vec{J}_1} \cos \phi_1 \right)$$

0 on resonance!

so, to lowest order:

$$\partial \langle H \rangle / \partial \vec{J}_1 = 0 \Leftrightarrow d\phi_1/dt = 0$$

is satisfied by resonance condition.

so $\vec{J}_{1,0}$ defined by resonance condition.

118

fixed points:

$$\hat{J}_{1,0} \leftrightarrow \text{resonant position} \\ \omega_1(\hat{J}) - S \omega_2(\hat{J}) = 0$$

$$\phi_{1,0} \leftrightarrow \sin \phi_1 = 0.$$

n.b.
see 22b

$$\begin{aligned} \langle H \rangle &= \langle H(\hat{J}_1, \hat{J}_2, \phi_1) \rangle \\ &= \langle H(\underbrace{\hat{J}_{1,0}}_{\text{resonance}} + \underbrace{\delta \hat{J}_1}_{\text{excursion}}, \phi_1 | \hat{J}_2) \rangle \end{aligned}$$

IOM

118, expanding:

$$\begin{aligned} \langle H(\hat{J}_1, \phi_1) \rangle &\approx H_0(\hat{J}_{1,0}) + \epsilon (H_{0,0}^{(4)}(\hat{J}_{1,0}) \\ &+ \cancel{\frac{\partial H_0}{\partial \hat{J}_1}} (\hat{J}_1 - \hat{J}_{1,0}) + \frac{\epsilon}{2} \frac{\partial^2 H_0}{\partial \hat{J}_1^2} (\hat{J}_1 - \hat{J}_{1,0})^2 \\ &+ 2 \epsilon H_{1,-5}^{(4)} \cos \phi_1 \end{aligned}$$

reson $\hat{J}_{1,0}$

⇒

$$\langle H(\hat{J}_1, \phi_1) \rangle \approx \text{const.} + \frac{\epsilon}{2} \frac{\partial^2 H_0}{\partial \hat{J}_1^2} (\hat{J}_1 - \hat{J}_{1,0})^2 + 2 \epsilon H_{1,-5}^{(4)} \cos \phi_1$$

ω_J have arrived at averaged Hamiltonian near resonance:

$$\langle H(\hat{J}_1, \phi_1) \rangle = \frac{1}{2} (\hat{J}_1 - \hat{J}_{1,0})^2 \left. \frac{\partial^2 H_0}{\partial \hat{J}_1^2} \right|_{\hat{J}_{1,0}} - F \cos \phi_1$$

$$= \frac{G}{2} (\hat{J}_1 - \hat{J}_{1,0})^2 - F \cos \phi_1$$

$$G = \left. \frac{\partial^2 H_0}{\partial \hat{J}_1^2} \right|_{\hat{J}_{1,0}}, \quad F = \overline{\epsilon} H_{1,0}^{(1)}$$

→ isomorphic to pendulum!

Recall for pendulum:

$$L = \frac{m l^2}{2} \dot{\theta}^2 - m g l (1 - \cos \theta)$$

$$H = p \dot{\theta} - L = \frac{p^2}{2 m l^2} - m g l \cos \theta$$

$$\Rightarrow H(\hat{J}_1, \phi) = \frac{\sigma}{2} (\hat{J}_1 - J_{1,0})^2 - F \cos \phi$$

is form of Hamiltonian near resonance.

Note:

- assumes $\frac{\partial^2 H}{\partial \hat{J}_1^2} = \frac{\partial \omega}{\partial \hat{J}_1} \neq 0$ (NL/shear)

"accidental" resonance.

- for properties:

$$\langle H(\hat{J}_1, \phi) \rangle = \frac{\sigma}{2} (\hat{J}_1 - J_{1,0})^2 - F \cos \phi$$

↓
shear/NL
parameter

↓
perturbation
amplitude

and so:

$$\begin{aligned} \Delta \dot{J} &= -F \sin \phi \\ \dot{\phi} &= \sigma \Delta J \end{aligned}$$

$$\begin{aligned} \phi &= 0 + \delta \phi \\ \Delta \dot{J} + F \sigma \delta \phi &= 0 \\ \text{near } \phi &= 0 \end{aligned}$$


e.e.

$$\Delta \dot{J}_1 = -F \cos \phi_{1,0} G \Delta J$$

$$\Delta \ddot{J}_1 + FG \cos \phi_{1,0} \Delta J = 0$$

$FG > 0 \Rightarrow \phi_1 = 0$, stable fixed point
(0-pt/elliptic point) ↗

$\phi_1 = \pm \pi \Rightarrow$ unstable fixed pt.
(x-pt/hyperbolic pt.)



→ stable fixed pt. \Leftrightarrow elliptic point \Leftrightarrow O pt.
 - island center



- center of trapped or libration region

→ unstable fixed point \Leftrightarrow hyperbolic point \Leftrightarrow X pt.

- island edge

- separatrix crossing point

→ separatrix ('separator') region of rotation (i.e. untrapped) from region of trapped (i.e. libration)



→ libration: elliptic orbits
 rotation: hyperbolic orbits

- width of separatrix - "island width"

$$\begin{aligned}
 \Delta J)_{\max} &\approx 2(E/G)^{1/2} \\
 &\approx 2 \left(-2G \frac{H_0 - S}{\omega} / \left. \frac{d^2 H}{dJ^2} \right|_{J_0} \right)^{1/2}
 \end{aligned}$$

i.e. particle + wave:


$$H = (p + m\omega/k)^2 / 2m + \sum \phi_0 \cos kx$$

$$\Delta p = (\sum \phi_0 \cdot m)^{1/2}$$

$$\Delta v \approx (\sum \phi_0 / m)^{1/2} \rightarrow \text{trapping width}$$

⇒ the Big Picture:

- resonant perturbations distort and foliate resonant tori in phase space, forming island chain structures.





Note:

- structure localized to resonant surface
- trapped } orbits stay { trapped
 untrapped } untrapped.
- resonant surface is foliated but not destroyed.
- motion remains on surface, though surface is ruffled...