

MHD Turbulence II

- Anisotropic Cascades and Critical Balance → A closer look.
- Extending the 4/5 Law.
- Selective Decay and Relaxation.
- 2D MHD - A Study in Turbulent Relaxation.

(a) Anisotropic Cascades and Critical Balance - A closer look.

Recall: I-k Phenomenology:

$$\epsilon \cong \frac{Z(l)^2}{T_{tr}(l)} \quad (\text{Bo weak, but RMS generated})$$

$$1/T_{tr}(l) \cong \frac{Z(l)^2}{l^2} T_A \quad T_A \sim \frac{l}{b_0 \text{ rms}}$$

$$\Rightarrow Z_e \sim (b_0 G)^{1/4} l^{1/4}$$

$$E(k) \sim \sqrt{\epsilon k_0} k^{-3/2}$$

and with strong Bo:

$$\epsilon \sim \frac{Z(l_{\perp})^2 Z(l_{\parallel}^2)}{\rho_{\perp}^2 |k_{\parallel} v_A|}$$

so W.T.T. Alfven's cascade:

$$E(k_{\perp}, k_{\parallel}) \sim (\epsilon V_A)^{1/2} / k_{\parallel}^{1/2} k_{\perp}^2 \sim \begin{matrix} \text{"hard"} \\ \text{in } k_{\perp} \end{matrix}$$

However, note:

$$\tau(k_{\perp}) \sim \delta B(k_{\perp}) \sim (\epsilon |k_{\parallel} V_A|)^{1/4} k_{\perp}^{1/2}$$

so

$$\delta B \cdot \tau_{\perp} \sim \frac{1}{k_{\perp}^{1/2}} (\epsilon |k_{\parallel} V_A|)^{1/4}$$

But recall:

- Alfven wave:

$$\omega \approx k_{\parallel} V_A$$

derived from:

$$\partial_t A = B_0 \nabla_{\parallel} \phi + \dots$$

$$\partial_t \nabla^2 \phi = B_0 \nabla_{\parallel} J_{\parallel} + \dots$$

$$\nabla_{\parallel} = \underbrace{\partial_z}_{\text{Linear}} + \frac{\delta B_{\perp}}{B_0} \nabla_{\perp}$$

Nonlinear

$$\text{Ratio } \frac{\text{Nonlinear}}{\text{Linear}} = k_{\text{cu}} = \frac{\partial B_{\perp} \cdot \sigma_{\perp}}{B_0} \downarrow \downarrow z$$

Kubo #

$$\Rightarrow k_{\text{cu}} \sim \frac{\partial B_{\perp} \text{ fac}}{B \Delta_{\perp}} \quad b_{\text{cu}} \rightarrow \text{parallel auto correlation length}$$

see stochastic fields discussion of Phys 235 2015.

$$\Delta_{\perp} \rightarrow \perp \text{ correlation length.}$$

Point: $B_0 \partial_z C + \partial B_{\perp} \cdot \sigma_{\perp} C = 0$

$$\partial_z C + \frac{\partial B_{\perp} \cdot \sigma_{\perp}}{B_0} C = 0$$

$$k_{\text{cu}} < 1 \rightarrow C \text{ evolves by many kicks in } \Delta_{\perp} \rightarrow \text{diffusion}$$

\rightarrow in WTT wave interactions are diffusive in character.

$$k_{\text{cu}} > 1 \rightarrow C \text{ scattered } > \Delta_{\perp} \text{ in one step}$$

\rightarrow fast transport in random media \rightarrow percolation

Analogy $\nabla_{\perp} C + \underline{v} \cdot \nabla C = 0$

$$k_{\perp} u = \frac{\nabla_{\perp} T_{ac}}{\Delta_{\perp}}$$

So we have a concern:

→ Physics of ~~the~~ MHD turbulence understood in terms of AlFven wave interactions.

→ but scalings of WTT spectrum suggest that wave character lost as cascade progresses

i.e.

$$k_{\perp} u \sim \frac{k_{\perp}^{-1}}{\rho_{\perp}^{1/2}} [E |k_{\perp} v_{\perp}|]^{1/4}$$

↑ as $\rho_{\perp} \downarrow$

i.e. How high can k_{\perp} # go and still be consistent with physics of AlFven Wave Cascade

⇒ Critical Balance Conjecture

(GS 75, KP '78)

⇒ MHD inertial range in strong field will set at $ku \sim 1$.

d.e. $\rightarrow dB_{\perp} \cdot v_{\perp} \sim \frac{Z(l_{\perp})}{l_{\perp}} \sim B_0 D_{\perp}$

so $\frac{Z(l_{\perp})}{l_{\perp}} \approx ku VA$

$\rightarrow \frac{T_A}{T_{\text{eddy}}} \rightarrow 1 \quad T_{Tr} \rightarrow T_{\text{eddy}} \sim T_A$

d.e. all timescales equalize
 $\rightarrow ku \sim 1$ is maximum ku and still retain Alfvénic character.
 \rightarrow why?

Recall:

- WTT $T_{T0} \sim T_{ac}$ } Triad coherence set by wave dispersion

$\rightarrow \pi \delta(\omega_{\underline{k}} - \omega_{\underline{k}'} - \omega_{\underline{k}''})$

- STT - Renormalized Theory

$T_{T0} \sim T_0$ } Triad coherence set by nonlinear scattering, etc.

$\rightarrow I / \Delta\omega_{\underline{k}} + \Delta\omega_{\underline{k}'} + \Delta\omega_{\underline{k}''}$
 $= \mathcal{O}_{\underline{k}, \underline{k}', \underline{k}''}$

So, renormalized wave interaction theory \Rightarrow

$$\Theta_{k, k', k''} = \frac{\Delta\omega_k + \Delta\omega_{k'} + \Delta\omega_{k''}}{(\omega_k - \omega_{k''} - \omega_{k'})^2 + (\Delta\omega_k + \Delta\omega_{k'} + \Delta\omega_{k''})^2}$$

\rightarrow recovers both limits \checkmark

Now, $\Theta_{k, k', k''}$ clearly sets T_{tr} .

So, can re-write phenomenological transfer balance as:

$$E \sim \frac{1}{l_\perp^2} \frac{Z(l_\perp)^2 Z(l_\perp)}{\sqrt{T_{tr}(l_\perp)}}$$

$$1/\sqrt{T_{tr}(l_\perp)} = \left[\underbrace{(k_{\perp} v_A)^2}_{\text{comparable to } k_{\perp} v_A} + \underbrace{\left(\frac{Z(l_\perp)}{l_\perp}\right)^2}_{\text{comparable to } k_{\perp} v_A} \right]^{1/2}$$

by analogy with $\Theta_{k, k', k''}$.

$$\textcircled{1} > \textcircled{2} \rightarrow \text{W.T.T.}$$

$$\textcircled{1} < \textcircled{2} \rightarrow \text{S.T.T.}$$

$$E \sim \frac{1}{l_{\perp}} \frac{Z(l_{\perp})^2 Z(l_{\perp})^2}{Z(l_{\perp})/l_{\perp}}$$

$$\sim Z(l_{\perp})^3 / l_{\perp}$$

and $Z(l_{\perp}) \sim (E l_{\perp})^{1/3}$

Point: $\langle Z(k)^2 \rangle \sim E^{2/3} k_{\perp}^{-5/3}$
but different physics!

- Back to
k 4)!

- GS spectrum.

- softer than
WTT.

- Great Power
Load of sky!

- $\frac{Z(l_{\perp})}{l_{\perp}}$ vs. $k_{\perp} V_A$

$$\frac{(E l_{\perp})^{1/3}}{l_{\perp}} \sim \frac{E^{1/3}}{l_{\perp}^{2/3}}$$

\rightarrow rate increases
as $l_{\perp} \downarrow$

\rightarrow contrast constant
 $\{k_{\perp} V_A\}$

$$- \frac{Z(l_{\perp})}{l_{\perp}} \sim \frac{E^{1/3}}{l_{\perp}^{2/3}} \rightarrow \frac{\delta B}{B_0} \cdot D_{\perp}$$

then $k_{\perp} \sim \pm \Rightarrow$

$$\frac{E^{1/3}}{k_{\perp}^{2/3}} \sim k_{\parallel}$$

$$\Rightarrow \boxed{k_{\parallel} \sim E^{1/3} k_{\perp}^{2/3}} \quad - \text{GS core.}$$

\rightarrow - Critical Balance is a hypothesis.

- Plausible answer to question of "how maintain Alfvénic cascade in state of strong (i.e. non-weak) turbulence?"

- anisotropy of spectrum supported by simulations (cf. Galtier).

BUT

- hypothesis, only.

\rightarrow Computational support semi-quantitative,

\rightarrow 5/3 vs 3/2 etc. still ongoing.

→ A word about triads.

In wave turbulence cascade, must satisfy:

$$\underline{k} = \underline{p} + \underline{q}$$

$$\omega_{\underline{k}} = \omega_{\underline{p}} \pm \omega_{\underline{q}} \quad (\text{WTT})$$

→ resonance criterion

Conditions satisfied by:

$$q_{\parallel} = 0$$

(i.e. \underline{q} is a cell,

so $\underline{k}_{\parallel} = p_{\parallel}$

driven by beats)

$$\underline{k}_{\perp} = \underline{p}_{\perp} + \underline{q}_{\perp}$$

and $\omega_{\underline{k}} = \omega_{\underline{p}} \pm \omega_{\underline{q}}$

⇒ - deformation of Alfvénic wave packet directly related to its interaction with 2D part of wave packet travelling in opposite direction.

- interaction passive w/r $\underline{k}_{\parallel}$
 ⇒ \perp transfer in long time limit.

- Inverse {cascades} of magnetic helicity would set up "selective decay" scenario

ie. magnetic energy scattered to small scale and dissipated
 \Rightarrow relaxation

magnetic helicity inverse cascades
 \Rightarrow avoids dissipation. Constraint; as survives.

c.f. {Frisch (75), Pouquet, et al. (76)
 (postscript)
 see also: Montgomery

- Why, where from?

\rightarrow Primarily: Statistical Mechanics


\rightarrow c.f.: Frisch '75, though not transparent.

easier \rightarrow "Taylor in Flatland" Problem.

Recall: Relaxation $\left\{ \begin{array}{l} \text{minimized } \langle B^2 \rangle \\ \text{conserving } \langle A^2 \rangle \end{array} \right.$

Does this follow from selective decay?

⇒ Explore Absolute Equilibrium



- remove forcing, dissipation etc.
- input excitations.

For 2D MHD (ignoring cross helicity):

have $A \Rightarrow X_i$
↳ mode amplitude

∴

$$E_M = \sum_{i=1}^N k_i^2 X_i^2$$

$$H = \sum_{i=1}^N X_i^2 \quad - \langle A^2 \rangle$$

$$\phi \rightarrow y_i$$

$$E_k = \sum_{i=1}^N k_i^2 y_i^2$$

Now, $H \rightarrow \alpha$ > conserved
 $E = E_M + E_k \rightarrow \beta$ > conserved

conserved, so PDF of this closed system is given by micro-canonical ensemble/distribution:

$$P(x, y) = \underset{\text{norm}}{C} \exp \left[- \sum_{i=1}^N \left[(\alpha + \beta k_i^2) x_i^2 + \beta k_i^2 y_i^2 \right] \right]$$

and can integrate out y_i (kE) part, so:

$$P(x) = C \exp \left[- \sum_{i=1}^N (\alpha + \beta k_i^2) x_i^2 \right]$$

then:


$$\begin{aligned} \langle A^2(k) \rangle &= \int dx_i x_i^2 P(x_i) \\ &= 1 / (\alpha + \beta k^2) \end{aligned}$$

$$\langle B^2(k) \rangle = \left[k^2 / (\alpha + \beta k^2) \right]$$

∞ observe immediately ∴



" A^2 wants remain at large scale"



" B^2 approaches equipartition"

⇒ A^2 distribution most populated at ~~smaller~~ large scales. Delay at small

⇒ B^2 distribution most populated at smaller. Approaches equipartition at small scale.

∴ - suggests A^2 populates large scales, B^2 approaches equipartition.

- suggestive of inverse cascade

of A^2 , along with forward cascade of energy.

- supports Selective Decay Hypothesis as foundation for "Taylor in Flatland".
- similar story ~~for~~ for Magnetic Helicity, though more laborious.

N.B. For 2D Fluid:

$$E = \int d^2x (\nabla\phi)^2 \quad - \text{energy}$$

$$\Omega = \int d^2y (\nabla^2\phi)^2 \quad - \text{enstrophy}$$

$$\Omega_i = k_i^2 E_i$$

$$\underline{v} \rightarrow \underline{X}_i$$

$$P(\underline{X}) = C \exp \left[- \sum_{i=1}^N (\alpha + \beta k_i^2) X_i^2 \right]$$

so
$$\langle v^2(k) \rangle = 1/(\alpha + \beta k^2)$$

$$\Omega(k) = k^2 / (\alpha + \beta k^2)$$

similar suggestion of dual cascade and minimum enstrophy state.

→ Is this story true?

⇒ What does dynamics tell us?
 Consider interactions in 2D MHD.

Observes:

- Reduced MHD

$$\frac{\partial \psi}{\partial t} + \frac{\nabla_{\perp} \phi \times \hat{z}}{\underline{1}} \cdot \frac{\nabla_{\perp} \psi}{\underline{1}} = B_0 \partial_z \phi + \eta \nabla_{\perp}^2 \psi$$

- 2D MHD

$$\frac{\partial \psi}{\partial t} + \nabla_{\perp} \phi \times \hat{z} \cdot \nabla_{\perp} \psi = \eta \nabla_{\perp}^2 \psi$$

so, with strong B_0 :

$$\langle A \cdot B \rangle \rightarrow \langle \psi \rangle B_0$$

so mean $\langle \psi \rangle$ in 2D captures magnetic helicity dynamics in strongly magnetized system.

For $\langle A^2 \rangle_{\perp}$ transfer, consider closure

of $\partial_t \langle A^2 \rangle$ equation, much akin to wave kinetics, though closure required.

See: Diamond, Hughes, Kim (posted).

Can write (see DHK) :

$$\frac{1}{2} \left[\partial_t \langle A^2 \rangle_{\underline{u}} + T(k) \right] = - \Gamma_A(k) \frac{\langle A \rangle}{\partial x} - \eta \langle B^2 \rangle_{\underline{u}}$$

triplet
 $\langle 0 \cdot \langle \nabla A^2 \rangle \rangle_{\underline{u}}$


flux

$$\Gamma_A = \left[\begin{matrix} \Gamma_0^{\phi}(\underline{u}) \langle \psi^2 \rangle_{\underline{u}} \\ - \Gamma_0^A(k) \langle B^2 \rangle_{\underline{u}} \end{matrix} \right]$$


$$\Gamma_{\underline{u}} = \sum_{\underline{u}} (\underline{u} \cdot \underline{u}' \times \underline{z})^2 \textcircled{1} \left\{ \langle \psi^2 \rangle_{\underline{u}'} \right. \\ \left. - \frac{(\underline{u}'^2 - k^2)}{|\underline{k} + \underline{u}'|^2} \langle A^2 \rangle_{\underline{u}'} \right\} \langle A^2 \rangle_{\underline{u}}$$

$$- \sum_{\substack{\underline{u} = \underline{u}' + \underline{z} \\ \underline{u}', \underline{z}}} (\underline{u} \cdot \underline{z} \times \underline{z}')^2 \textcircled{2} \langle A^2 \rangle_{\underline{u}'} \langle \psi^2 \rangle_{\underline{z}}$$

- ①, ③ → coherent damping, incoherent emission
- akin to scattering of passive scalar, → small scale / chop-up.
- conserve $\langle \psi^2 \rangle$ upon $\sum_{\underline{u}}$ together.



- ② → coherent damping/growth - from back reaction (J x B) into Ohm's law.
- reshuffle $\langle A^2 \rangle$ to larger scale. Sign k'^2 vs k^2 !
- \sum_n conserved ~~quantities~~ $\langle A^2 \rangle$ independently.



→ correspondence to condensation of water (currents) attracting.

- ① + ② → net effective reactivity sign.
- see Γ_A , too. - 'negative reactivity'
- Alfvénized state

$\Rightarrow E_k > E_M \Rightarrow$ ~~shuffled~~ $\langle A^2 \rangle_n$ shuffled to smaller scale.

$E_M < E_k \Rightarrow \langle A^2 \rangle_n$ transferred to larger scale.

and transfer need not be local.

⇒ In dynamics ~~the~~ $\langle A^2 \rangle$; $\langle A \cdot B \rangle$
evolution is complex.

⇒ N.B. Recall Flux expulsion:

$$\frac{V_{A0}^2}{V^2} R_m < 1 \rightarrow A \text{ @ } \rho \text{-surface } B \text{ expelled}$$

$$> 1 \rightarrow J \times B \text{ disrupts vortex, expulsion}$$

$$\Rightarrow B_0^2 < \rho_0 V^2 / R_m$$

but $\langle \tilde{B}^2 \rangle \gg B_0^2$, upon stretching,
weak B_0 is sufficient!

Zeldovich:

$$\frac{\partial A}{\partial t} + \underline{v} \cdot \nabla A = -v_r \frac{\partial \langle A \rangle}{\partial x} + \eta \nabla^2 A$$

*A and avg. ⇒

$$\eta \langle \tilde{B}^2 \rangle = \langle \tilde{v}_r \tilde{A} \rangle \frac{\partial \langle A \rangle}{\partial x}$$

$$\langle \tilde{B}^2 \rangle = \frac{\eta_r}{\eta} B_0^2$$

$$= \frac{\tilde{v} \tilde{v}_r}{\eta} B_0^2 \approx R_m B_0^2 \checkmark$$

so, crudely:

$$\langle B^3 \rangle / R_m > \langle \rho V^2 \rangle / R_m$$

✓

⇒ Questions still open ↓

∴ Taylor conjecture remains a conjecture ↓